

Provided for non-commercial research and educational use only.
Not for reproduction or distribution or commercial use.



This article was originally published in a journal published by Elsevier, and the attached copy is provided by Elsevier for the author's benefit and for the benefit of the author's institution, for non-commercial research and educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are prohibited. For exceptions, permission may be sought for such use through Elsevier's permissions site at:

<http://www.elsevier.com/locate/permissionusematerial>

Research paper

The intraspleen huPBL NOD/SCID model to study the human HIV-specific antibody response selected in the course of natural infection

Sophia Steyaert^a, Lieven Verhoye^a, Els Beirnaert^{b,1}, Helen Donners^b,
Katrien Fransen^b, Leo Heyndrickx^b, Guido Vanham^{b,c},
Geert Leroux-Roels^a, Peter Vanlandschoot^{a,*}

^a Ghent University and Hospital, Department of Clinical Chemistry, Microbiology and Immunology, Centre for Vaccinology, De Pintelaan 185, 9000 Ghent, Belgium

^b Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium

^c Department of Biomedical Sciences, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

Received 4 September 2006; received in revised form 23 November 2006; accepted 1 December 2006

Available online 9 January 2007

Abstract

The intrasplenic injection of human peripheral blood mononuclear cells (PBMCs) into severely immune deficient NOD/SCID mice, causes a massive and transient dominant expansion of human B cells in the spleen. This permits the easy isolation of human monoclonal antibodies specific for different antigens by a Kohler and Milstein-based method. Here we studied the human HIV-specific antibody response in the circulation of mice after intrasplenic transfer of PBMC from untreated HIV-infected patients with detectable to high viral load as well as from HAART-treated and from untreated patients, who kept an undetectable viral load (the latter referred to as “natural suppressors”). Excellent B cell expansion was obtained for all PBMC. High level replication of virus was observed after transfer of PBMC of untreated viremic patients only. A strong and multispecific HIV-specific antibody response was observed after transfer of PBMC of untreated viremic patients and natural suppressors. In contrast, only a weak and pauci-specific antibody response was detected in mice reconstituted with PBMC from successfully treated patients. Based on these observations we conclude that the use of the intraspleen mouse model confirmed a) the presence of HIV-specific circulating memory B cells in untreated patients and natural suppressors; b) the nearly complete absence of circulating memory B cells in patients receiving highly active antiretroviral therapy. Using the intraspleen model we generated large numbers of immortalized B cells and isolated two anti-p24 human monoclonal antibodies. We further conclude that the intraspleen huPBL NOD/SCID model is a small animal model useful for the analysis of the antibody response against HIV found in patients. © 2006 Elsevier B.V. All rights reserved.

Keywords: NOD/SCID; HIV; Human; Hybridoma; Monoclonal antibody

Abbreviations: HAART, highly active antiretroviral therapy; HCV, hepatitis C virus; HIV, human immunodeficiency virus; huPBL, human peripheral blood lymphocyte; NOD/SCID, non-obese diabetic/severely combined immunodeficiency; PBMC, peripheral blood mononuclear cell.

* Corresponding author. Ghent University and Hospital, Centre for Vaccinology, Building A, 1st floor, De Pintelaan 185, 9000 Ghent, Belgium. Tel.: +32 9 240 60 39; fax: +32 9 240 63 11.

E-mail address: Peter.Vanlandschoot@UGent.be (P. Vanlandschoot).

¹ Present address: Ablynx Technologiepark, 9052 Zwijnaarde.

1. Introduction

Immune deficient mice in which human cells survive and function are a valuable tool for the *in vivo* study of human immune function and human diseases including auto-immune disorders, lymphoma and infectious diseases (Murphy et al., 1996). In studies of HIV pathogenesis and therapy, the two most commonly used systems are the SCID-hu model and the huPBL-SCID model (McCune et al., 1988; Mosier et al., 1988; Borkow, 2005). In the latter model, peripheral blood mononuclear cells (PBMC) are commonly injected into the peritoneum of the severely immune deficient mice resulting in a human cell graft in which T cells predominate. We described a variant huPBL NOD/SCID model, in which PBMC were transferred into the spleen instead of into the peritoneum (Depraetere et al., 2001). In brief, this intraspleen (*i.s.*) transfer results in a massive non-specific, T cell-dependent expansion of B cells that actively secrete immunoglobulins in the blood of the mice. The antigen specificities of the mouse sera reflect the specificities found in the human B cell donor, elicited by infections or immunizations. Six to ten days post injection, human B cells are transiently predominant in the mouse spleen from where they can be easily recovered for subsequent immortalization procedures and generation of human monoclonal antibodies. So far we have successfully generated human monoclonal antibodies against the hepatitis B surface and core antigens (Cao et al., 2001; Vanlandschoot et al., 2002), hepatitis C NS3A and NS4B1 (Depraetere et al., 2001) and tetanus toxoid (unpublished results). We envisage that the *i.s.* huPBL NOD/SCID model may be used to isolate therapeutic human antibodies and to analyse the antibody repertoire in immune and vaccinated individuals to help in the design of vaccines against viruses like HCV and HIV (analytic vaccinology).

In the present study we assessed whether *i.s.* transfer of PBMC from HIV-1 infected patients would result in a good B cell expansion, circulation of HIV-specific antibodies and allow for the generation of monoclonal HIV-specific antibodies. Success is not necessarily guaranteed since HIV infection affects both number and function of T and B lymphocytes. CD4⁺T lymphocytes drive the expansion of B cells after *i.s.* transfer of PBMC in mice (Depraetere et al., 2001). However, HIV infection is known to cause a rapid depletion of human CD4 T cells in SCID mice (Mosier et al., 1991). B cell behaviour is altered as represented by a paradoxical polyclonal activation of B cells (Lane et al., 1983), resulting in an increased frequency of circulating antibody secreting cells and hypergamma-

globulinemia (Yarchoan et al., 1986; Martinez-Maza et al., 1987; Amadori et al., 1988; Amadori et al., 1989; Shirai et al., 1992). During untreated and HAART-treated chronic HIV infection, memory B cells gradually disappear from the circulation (De Milito et al., 2001; Fondere et al., 2003; Chong et al., 2004; Titanji et al., 2006). A deregulated expression of several surface molecules (Indraccolo et al., 1993; Samuelsson et al., 1997; Moir et al., 2001; Moir et al., 2004) and a decreased response to B cell activators *in vivo* and *in vitro* have been described (Lane et al., 1983; Conge et al., 1998).

2. Materials and methods

2.1. Human samples

Informed consent was obtained from all participants and the study protocol was approved by the ethical committees of the Ghent University Hospital, Belgium (project number 2002/109) and the Institute of Tropical Medicine (reference number 01 10 1 373).

EDTA anti-coagulated blood was drawn from HIV-1 infected patients attending the HIV outpatient clinic either at the Ghent University Hospital (CV) or at the Institute of Tropical Medicine (VI). Plasma was kept frozen at -80°C until use. PBMC were prepared by Ficoll–Hypaque centrifugation (density 1.077 g/ml) (Nycomed Pharma) and stored in liquid nitrogen. Aliquots of these PBMC were used to produce the chimeric mice.

Table 1
Patient characteristics at the time of blood sampling

	Subject	HAART ^a	Viral load (log geq/ml)	CD4 (#/μl)
Group A	VI 2677	–	5.58	269
	VI 2592	–	5.28	378
	VI 2586	–	4.62	595
	CV7	–	3.92	1090
	CV8	–	3.92	345
	VI 2588	–	2.38	612
Group B	VI 2587	NRT ^b +PI ^c	<1.7	979
	VI 2589	NRT+PI	<1.7	592
	VI 2571	NRT+NNRT ^d	<1.7	631
	VI 2668	NRT+NNRT	<1.7	809
	VI 2670	NRT+PI	<1.7	678
	VI 2675	NRT+NNRT	<1.7	609
Group C	CV16	–	<1.7	663
	CV25	–	<1.7	810
	CV28	–	<1.7	846

^a HAART highly active antiretroviral therapy.

^b NRT nucleoside reverse transcriptase inhibitor.

^c PI protease inhibitor.

^d NNRT non-nucleoside reverse transcriptase inhibitor.

2.2. huPBL NOD/SCID mice

NOD/LtSz-Prkdc^{scid}/Prkdc^{scid} (NOD/SCID) mice are severely immune deficient, since they lack functional B and T cells, and display reduced innate immunity and haemolytic complement (Shultz et al., 1995). These mice were bred and maintained under specific pathogen-free conditions. The mice were housed in individually ventilated cages and fed with autoclaved food and water. Mice were used between eight and twelve weeks of age. The study was approved by the animal ethical committee of the Faculty of Medicine, University Ghent, Belgium (ECP 01/16). To generate chimeric animals, NOD/SCID mice were conditioned by the injection of 0.5 mg TM β 1, a rat monoclonal antibody directed against the murine IL-2

receptor β -chain (Tanaka et al., 1991; Tanaka et al., 1993), and total body gamma irradiation (3 Gy) generated by a linear accelerator. Twenty-four hours later, 5 to 10 $\times 10^6$ PBMC from HIV-1 infected patients were injected i.s. For fusion experiments, 2 $\times 10^7$ cells were injected i.s. PBMC from each patient were injected into three mice.

2.3. Quantification of human IgG and detection of HIV-specific antibodies

Total human IgG concentrations were determined in mouse plasma using an in-house ELISA as described previously (Tournoy et al., 2000).

HIV-specific antibody profiles were determined in human and mouse plasma using INNO-LiA HIV




Fig. 1. Follow-up details for the three subjects in group C (CV16, CV25 and CV28). Viral loads are represented by black triangles and CD4 absolute counts by grey circles. AZT, zidovudine.

confirmation strips (Innogenetics). Each strip is coated with different antigens of HIV-1 (gp120, gp41, p31, p24 and p17) and HIV-2. Antibodies against these antigens are detected by the development of a line which is scored between 0 and 4 according to the manufacturer's instructions. Samples were diluted 1/100 according to the manufacturer's instructions.

Hybridoma supernatant was screened for HIV-specific antibodies using the Ortho HIV-1/HIV-2 Ab-capture ELISA Test System.

2.4. Determination of viral load

Viral load was measured using COBAS Amplicor HIV-1 Monitor™ version 1.5 (Roche) according to the manufacturer's instructions. The limit of detection for human samples was 1.7 log geq/ml and for mouse samples 3.6 log geq/ml. The mouse samples were diluted due to the limited sample volume available.

2.5. Hybridoma techniques

Spleen cells from the reconstituted NOD/SCID mice and K6H6/B5 heteromyeloma cells were washed in calcium-free phosphate buffered saline (PBS) and mixed at a 3:1 ratio. Fifty percent polyethylene glycol (PEG) 1500 (Boehringer Mannheim) was added for 2 min and washed away. Fused cells were cultured in medium supplemented with 10 µg/ml human recombinant insulin (Boehringer Mannheim), 10% Fetal Clone I serum (HyClone), and 10% BM Condimed H1 (Boehringer Mannheim). Irradiated PBMC from a healthy seronegative donor were added as feeder cells. Fusion medium containing hypoxanthine-aminopterin-thymidine (HAT) (Life Technologies), and ouabain (1 µM; Sigma) was added the day after the fusion. The K6H6/B5 heteromyeloma cell line is HAT sensitive and ouabain resistant. Cultures were replenished with fresh medium every three days.

2.6. Statistical analysis

Statistics were performed using Mann–Witney *U*-test and Kruskal–Wallis test (Medcalc program version 7.1.0.1) (Schoonjans et al., 1995).

3. Results

3.1. Successful expansion of B cells after i.s. injection of PBMC from HIV-1 infected patients

PBMC were isolated from the blood of 15 HIV-1 infected patients who were divided into three groups

according to their treatment status and the viral burden in their plasma (Table 1). Group A consisted of six individuals who did not receive HAART for at least six months and whose plasma viral load ranged from 2.38 to more than 5.88 log geq/ml. The second group (group B) consisted of six subjects who currently received HAART and who kept their viral load below the limit of detection (<1.70 log geq/ml). The third group (group C) consisted of three subjects who did not receive antiviral agents and who naturally controlled their viral replication below the limit of detection (<1.70 log geq/ml). Longitudinal details of viral load and CD4 counts in these three natural suppressors are shown in Fig. 1.

Approximately 1×10^7 of these PBMC were used to reconstitute optimally conditioned NOD/SCID mice by direct injection in the spleen. Two out of 45 mice died before 7 days after transfer. Thirty mice survived until 14 days after transfer and only seven of these until one week thereafter. Death of the mice was due to a severe graft-versus-host disease which is easily diagnosed by the presence of a hunched back, ruffled fur, emaciation and loose stools. Excellent expansion of B cells was achieved in all mice (100%) as demonstrated by increasing human IgG levels in their blood (Table 2). The average IgG level of group A was 316 ± 64 µg/ml (mean \pm SEM) one week after injection and 4261 ± 1317 µg/ml two weeks after injection. For group B, these levels were 245 ± 44 µg/ml after one week and 2599 ± 437 µg/ml after two weeks. For group C, the average human IgG level was 291 ± 92 µg/ml after one week and 2871 ± 377 µg/ml after two weeks. These polyclonal human IgG levels did not differ significantly between groups ($p > 0.05$).

Table 2

Human polyclonal IgG production and viral load in mouse blood after i.s. transfer of PBMC from HIV-1 infected subjects

	Group A	Group B	Group C
<i>hIgG</i> (µg/ml)			
d7	316 \pm 64	245 \pm 44	291 \pm 92
d14	4261 \pm 1317	2599 \pm 437	2871 \pm 377
<i>Viral load</i> (log geq/ml)			
d7	4.99 \pm 0.24	<3.70 \pm 0.00*	<3.70 \pm 0.00*
d14	6.98 \pm 0.29	3.87 \pm 0.09*	<3.70 \pm 0.00*

Group A subjects did not receive antiretroviral therapy and their viral loads were above the limit of detection. Group B subjects received highly active antiretroviral therapy (HAART) and suppressed viral replication below the limit of detection. Group C subjects suppressed viral replication below the limit of detection without receiving antiretroviral therapy. Data are presented as mean \pm standard error of the mean (SEM). *statistically significant difference ($p < 0.01$) versus group A.




Fig. 2. HIV-specific antibody response in mouse blood one week and two weeks after i.s. transfer of PBMC from HIV-1 infected subjects and in human plasma. For mouse blood, the results are presented as a cumulative INNO-LiA score for the three mice per patient. For human plasma, the INNO-LiA score is shown for each patient. Group A is represented by black bars, group B by white bars and group C by grey bars. Statistical analysis was performed using Kruskal–Wallis test.

3.2. High viral burden after i.s. transfer of PBMC from patients with detectable viral load

As early as 7 days after reconstitution, high viral genome concentrations were measured in the blood of the mice reconstituted with PBMC from untreated viremic HIV patients (group A). These levels further increased one week later (Table 2). In contrast, viral replication was weak to absent in the blood of mice reconstituted with PBMC from treated patients and natural suppressors (group B and group C). The differences in viral replication between group A (4.99 ± 0.24 log geq/ml after one week and 6.98 ± 0.29 log geq/ml after two weeks) and group B (<3.70 log geq/ml after one week and 3.87 ± 0.09 log geq/ml after two weeks) and between group A and group C (<3.70 log geq/ml after one week and <3.70 log geq/ml after two weeks) were highly significant ($p < 0.01$). There was no correlation between the viral burden in mouse plasma and human IgG levels in mouse plasma ($p > 0.05$).

3.3. Induction of a strong and multispecific HIV-specific antibody response after i.s. transfer of PBMC from untreated patients and natural suppressors

As shown in Fig. 2, the HIV-specific antibody profile was strong and broadly reactive (gp120, p41, p31, p24 and p17) when PBMC from untreated viremic patients (group A) and natural suppressors (group C) were used. However, the HIV-specific reactivity was weak to absent and pauci-specific (gp120, gp41 and p24) with PBMC from patients who were receiving antiviral therapy (group B). Kruskal–Wallis analysis showed statistically significant differences at a p -level of < 0.1 for all antigens two weeks post transfer. For the untreated

patients (group A), the HIV-specific antibody response was more envelope directed (gp120 and gp41) and for the natural suppressors (group C), this response was more gag-directed (p24 and p17). In contrast with the antibody profile induced after transfer of PBMC in the mice, the HIV-specific antibody response in human plasma did not differ between groups ($p > 0.1$).

3.4. Immortalization of human HIV-specific B cells after i.s. transfer in NOD/SCID mice

As shown in Table 3, a stable hybridoma that produced human monoclonal antibodies against the HIV gag protein p24 was generated twice from PBMC of an untreated HIV-infected subject (CV17) whose viral load was very low (1.80 log geq/ml) and whose plasma antibodies did not show any broadly neutralizing activity (Steyaert et al., in press). The activated human B cells were recovered from the mouse spleen one week after injection. Previous experiments showed that at that time, human B cells are predominantly present in the spleen. The human B cells in experiment I were fused with the heteromyeloma cell line and then extensively diluted before seeding the wells at a frequency of 100 B cells per well (40×96 -well). After 21 days, clusters of viable cells were detected in 284 of 3840 seeded wells (7%). Although IgG secretion was detected in all of them, HIV-specific antibodies were only found in one. The cells were successfully subcloned until monoclonality (anti-p24) was obtained. In a second experiment (II), seeding of B cells was done at varying numbers: 10×96 wells were seeded at 100 B cells per well, 7×96 -well at 1000 B cells per well, 5×48 -well at 10,000 B cells per well and 2×6 -well at 100,000 B cells per well. In the wells seeded at the highest densities (100,000 and 10,000 B cells per well), clusters of immortalized B cells

Table 3
Summary of clonal yields and immortalization efficiencies for generation of human monoclonal antibodies to HIV-1 proteins

No of wells	# B cells/well	# of wells with growth	# wells producing IgG	# wells producing anti-HIV	# stable monoclonal anti-HIV ^a
<i>Exp I</i>					
40×96 well (100) ^b	100	284 (7)	284 (7)	1 (0.03)	1 (anti-p24) ^c
<i>Exp II</i>					
2×6 well (100)	100,000	12 (100)	ND ^d	1 (8)	0
5×48 well (100)	10,000	240 (100)	ND	32 (13)	0
7×96 well (100)	1000	372 (55)	ND	47 (7)	1 (anti-p24)
10×96 well (100)	100	38 (4)	ND	2 (0.21)	0

^a Anti-HIV production in approximately 70% of wells seeded at 1 cell/well for two consecutive rounds.

^b Values in parentheses represent percentages.

^c Specificity of the monoclonal antibody.

^d ND not done.

were obtained in all of them (100%). HIV-specific antibodies were present in 1 of 12 (8%) and in 32 of 240 (13%), respectively. However, this anti-HIV positivity decreased rapidly and all limited dilution steps were unsuccessful. In the wells seeded at the lowest concentrations (1000 to 100 B cells per well), viable cell clusters were present in 55% and 4% of wells respectively and anti-HIV antibodies in 47 of 372 (13%) and 2 of 38 (5%) wells, respectively. Again, a striking loss of anti-HIV positivity was observed. Out of the wells seeded at 1,000 B cells per well, one stable hybridoma cell line producing anti-p24 could be generated.

4. Discussion

This report describes the exploration of the i.s. huPBL NOD/SCID model (Depraetere et al., 2001) for transfer of PBMC from HIV-infected patients. The engraftment after i.s. transfer was excellent in all mice (100%). The functionality of the engrafted human B cells was demonstrated by an early and vigorous antibody response in the blood of the mice. The transfer of PBMC from untreated HIV-infected patients with detectable viremia in optimally conditioned NOD/SCID mice resulted in replication of endogenous virus. In the mice reconstituted with PBMC from patients having undetectable viral load, replication often remained below detection ($<3.60 \log \text{ geq/ml}$). We assume that the proviral load in the transferred PBMC was strongly related to the viral load in the plasma of the patients (Andreoni et al., 2000; Ngo-Giang-Huong et al., 2001).

The HIV-specific antibody response after transfer of PBMC from untreated viremic patients and natural suppressors was strong and multispecific (gp120, gp41, p31, p24 and p17). HIV-specific antibody responses were weak and pauci-specific (gp120, gp41 and p24) after transfer of PBMC from patients receiving HAART. Since the human circulating memory B cells constitute the basis for the antibodies present in the blood of the mice, we conclude that the treated HIV-infected patients lost most of the HIV-specific memory B cells in their circulation. While a gradual loss of circulating memory B cells during chronic HIV infection occurs (Titanji et al., 2006), it has indeed been reported that memory HIV-1 specific B cells disappear from the blood after 8 to 46 months of successful antiretroviral therapy (Fondere et al., 2003).

We also noted that the HIV-specific antibody profile after transfer of PBMC of untreated viremic patients differed from the HIV-specific antibody profile obtained after transfer of PBMC of natural suppressors. In the latter

group the antibody response was mainly directed against proteins encoded by the gag gene (p24 and p17) while the response of PBMC from the viremic subjects was mainly directed to the viral envelope (gp120 and gp41). Antibodies against env and gag antigens are mounted during an HIV infection but differ in the extent to which they are sustained throughout the course of an HIV infection. While anti-env is usually present at high titres even in individuals who progress to disease, the anti-gag response is lost during disease progression (Schmidt et al., 1989; Cheingsong-Popov et al., 1991; Zwart et al., 1994; Hogervorst et al., 1995; Strathdee et al., 1995; Binley et al., 1997). This decline in anti-gag is thought to be a surrogate marker for the loss in T cell help (Binley et al., 1997; Trkola et al., 2004). Therefore, it is most likely that the gag-directed antibodies in the untreated naturally controlling patients reflect good functional T cell help.

The HIV antibody profile differed between groups in the intrasplenic expanded PBMC. This divergence was however not present in the corresponding human plasma. This is most likely a result of compartmentalization. In the mouse model, antibody production is derived from the memory B cells present in the circulation of HIV-infected subjects. In contrast, the HIV-specific antibodies detected in human plasma are predominantly derived from plasma cells in the bone marrow and antibody producing cells in the lymph nodes (Morris et al., 1998). These antibodies have also been secreted over an extended period of time.

In order to isolate HIV-specific monoclonal antibodies using the i.s. huPBL NOD/SCID model, it is clear that PBMC from untreated patients need to be used because only such patients have HIV-specific memory B cells in their peripheral blood. We were able to generate large numbers of IgG producing hybridoma cells, but the isolation of monoclonal hybridoma cell lines by limited dilutions was more difficult. There are several possible explanations for this low success rate. First of all, untreated HIV patients have a reduced number of memory B cells. Secondly, it has been estimated that in untreated patients about 10^{-2} to 10^{-3} of the memory B cells are HIV-specific (Fondere et al., 2004). We previously showed that 7 days after injection of 2×10^7 PBMC, $\sim 8 \times 10^6$ human B cells are present in the spleen (Depraetere et al., 2001). This would suggest that only 8000–80,000 HIV-specific B cells are available for immortalization. Finally, we noticed that the immortalized B cells grew slowly and anti-HIV production was easily lost. Clearly, B cell function and response is deregulated by HIV infection and apparently HIV also affects the immortalized B cells. The slow growth of fused B cells and the loss of HIV-specific antibody production has also been observed by other authors (Buchacher et al., 1994). It

was hypothesized that genomic instability of the cells within the first weeks after immortalization was responsible for this problem. Nevertheless, we were able to isolate two stable human monoclonal cell lines, that produce an antibody against the gag protein p24. Human monoclonal antibodies with this specificity are mainly used for diagnostic purposes. So far, human monoclonal antibodies against HIV have been produced by immortalizing B cells with EBV (Gorny et al., 1989) or by fusing B cells with appropriate myeloma cells (Buchacher et al., 1994). These methods have however a very low efficiency and therefore, alternative strategies are to be developed. In our mouse model, PBMC are injected directly into the spleen. This results in a strong polyclonal activation and a striking expansion of B cells from the memory B cells present in the PBMC mix. Recently, such memory B cells have been used for the successful generation of human monoclonal antibodies against the SARS coronavirus by culturing them in the presence of EBV and suitable growth factors (Traggi et al., 2004). Whether this approach is successful for the generation of human monoclonal antibodies against HIV remains to be determined.

In conclusion, transplantation of human lymphoid cells from HIV-infected patients into the spleen of immune deficient NOD/SCID mice is feasible for the study of HIV-specific human humoral immune responses and for the isolation of human monoclonal antibodies against HIV. Intrasplenic transfer might also be used to increase the number of HIV-specific memory B cells, before a single-chain phage display library is constructed. The i.s. huPBL NOD/SCID model may therefore be helpful to identify and isolate monoclonal antibodies that bind broadly neutralizing, enhancing and non-neutralizing epitopes on HIV. Such antibodies will be useful for the development of candidate vaccines.

Acknowledgement

We thank Filip Van Wanzele (UZGent), Beatrijs Vandergucht (UZGent), Eric Florence (ITM) en Marc Vandenbruaene (ITM) for recruiting HIV-infected patients; Tom Boterberg (UZGent) for assistance in the irradiation of the animals; Marianne Mangelschots (ITM) for viral load determinations and Guido van der Groen (ITM) for early contributions. Research was supported by the Fund for Scientific Research, Flanders, Belgium (FWO-Vlaanderen, grant number G.0346.03) and the Concerted Research Initiative of the University of Ghent, Belgium (GOA, project number 12050203). S.S. was a research fellow of the Fund for Scientific Research, Flanders, Belgium (FWO-Vlaanderen).

References

- Amadori, A., DeRossi, A., Faulkner-Valle, G.P., Chieco-Bianchi, L., 1988. Spontaneous in vitro production of virus-specific antibody by lymphocytes from HIV-infected subjects. *Clin. Immunol. Immunopathol.* 46, 342.
- Amadori, A., Zamarch, R., Ciminale, V., Del Mistro, A., Siervo, S., Alberti, A., Colombatti, M., Chieco-Bianchi, L., 1989. HIV-1 specific B cell activation. A major constituent of spontaneous B cell activation during HIV-1 infection. *J. Immunol.* 143, 2146.
- Andreoni, M., Parisi, S.G., Sarmati, L., Nicastrì, E., Ercoli, L., Mancino, G., Sotgiu, G., Mannazzu, M., Trevenzoli, M., Tridente, G., Concia, E., Aceti, A., 2000. Cellular proviral HIV-DNA decline and viral isolation in naive subjects with <5000 copies/ml of HIV-RNA and >500×10⁶/l CD4 cells treated with highly active antiretroviral therapy. *AIDS* 14, 23.
- Binley, J.M., Klasse, P.J., Cao, Y., Jones, I., Markowitz, M., Ho, D.D., Moore, J.P., 1997. Differential regulation of the antibody responses to Gag and Env proteins of human immunodeficiency virus type 1. *J. Virol.* 71, 2799.
- Borkow, G., 2005. Mouse models for HIV-1 infection. *IUBMB Life* 57, 819.
- Buchacher, A., Predl, R., Strutzenberger, K., Steinfellner, W., Trkola, A., Purtscher, M., Gruber, G., Tauer, C., Steindl, F., Jungbauer, A., Katinger, H., 1994. Generation of human monoclonal antibodies against HIV-1 proteins; electrofusion and Epstein-Barr virus transformation for peripheral blood lymphocyte immortalization. *AIDS Res. Hum. Retrovir.* 10, 359.
- Cao, T., Lazdina, U., Desombere, I., Vanlandschoot, P., Milich, D.R., Sällberg, M., Leroux-Roels, G., 2001. Hepatitis B virus core antigen binds and activates naive human B cells in vivo: Studies with a human PBL-NOD/SCID mouse model. *J. Virol.* 75, 6359.
- Chengsong-Popov, R., Panagiotidi, C., Bowcock, S., Aronstam, A., Wadsworth, J., Weber, J., 1991. Relation between humoral responses to HIV gag and env proteins at seroconversion and clinical outcome of HIV infection. *BMJ* 302, 23.
- Chong, Y., Ikematsu, H., Kikuchi, K., Yamamoto, M., Murata, M., Nishimura, M., Nabeshima, S., Kashiwagi, S., Hayashi, J., 2004. Selective CD27⁺ (memory) B cell reduction and characteristic Bcell alteration in drug-naive and HAART-treated HIV type 1-infected patients. *AIDS Res. Hum. Retrovir.* 20, 219.
- Conge, A.-M., Tarte, K., Reynes, J., Segondy, M., Gerfaux, J., Zembala, M., Vendrell, J.-P., 1998. Impairment of B-lymphocyte differentiation induced by dual triggering of the B-cell antigen receptor and CD40 in advanced HIV-1-disease. *AIDS* 12, 1437.
- De Milito, A., Morch, C., Sonnerborg, A., Chiodi, F., 2001. Loss of memory (CD27) B lymphocytes in HIV-1 infection. *AIDS* 15, 957.
- Depraetere, S., Verhoye, L., Leclercq, G., Leroux-Roels, G., 2001. Human B cell growth and differentiation in the spleen of immunodeficient mice. *J. Immunol.* 166, 2929.
- Fondere, J.-M., Huguët, M.-F., Yssel, H., Baillat, V., Reynes, J., van de Perr, P., Vendrell, J.-P., 2003. Detection of peripheral HIV-1-specific memory B cells in patients untreated or receiving highly active antiretroviral therapy. *AIDS* 17, 2323.
- Fondere, J.-M., Huguët, M.-F., Macura-Biegun, A., Baillat, V., Ohayon, V., Reynes, J., Vendrell, J.-P., 2004. Detection and enumeration of circulating HIV-1-specific Memory B cells in HIV-1-infected patients. *J. Acquir. Immune Defic. Syndr.* 35, 114.
- Gorny, M.K., Gianakakos, V., Sharpe, S., Zolla-Pazner, S., 1989. Generation of human monoclonal antibodies to human immunodeficiency virus. *Proc. Natl. Acad. Sci. U. S. A.* 86, 1624.

- Hogervorst, E., Jurriaans, S., de Wolf, F., van Wijk, A., Wiersma, A., Valk, M., Roos, M., van Genem, B., Coutinho, R., Miedema, F., Goudsmit, J., 1995. Predictors for non- and slow progression in human immunodeficiency virus (HIV) type 1 infection: low viral RNA copy numbers in serum and maintenance of high HIV-1 p24-specific but not V3-specific antibody levels. *J. Infect. Dis.* 171, 811.
- Indraccolo, S., Mion, M., Zamarchi, R., Veronesi, A., Veronese, M.L., Panozzo, M., Betterle, C., Barelli, A., Borri, A., Amadori, A., Chicco-Bianchi, L., 1993. B cell activation and human immunodeficiency virus infection. V. Phenotypic and functional alterations in CD5+ and CD5- B cell subsets. *J. Clin. Immunol.* 13, 381.
- Lane, H.C., Masur, H., Edgar, L.C., Whalen, G., Rook, A.H., Fauci, A.S., 1983. Abnormalities of B-cell activation and immunoregulation in patients with the acquired immunodeficiency syndrome. *N. Engl. J. Med.* 309, 453.
- Martinez-Maza, O., Crabb, E., Mitsuyasu, R.T., Fahey, J.L., Giorgi, J.V., 1987. Infection with the human immunodeficiency virus (HIV) is associated with an in vivo increase in B lymphocyte activation and immaturity. *J. Immunol.* 138, 3720.
- McCune, J.M., Namikawa, R., Kaneshima, H., Shultz, L.D., Lieberman, M., Weissman, I.L., 1988. The SCID-hu mouse: murine model for the analysis of human hematolymphoid differentiation and function. *Science* 241, 1632.
- Moir, S., Malaspina, A., Ogwaro, K.M., Donoghue, E.T., Hallahan, C.W., Ehler, L.A., Liu, S.Y., Adelsberger, J., Lapointe, R., Hwu, P., Baseler, M., Orenstein, J.M., Chun, T.W., Mican, J.A.M., Fauci, A.S., 2001. HIV-1 induces phenotypic and functional perturbations of B cells in chronically infected individuals. *Proc. Natl. Acad. Sci. U. S. A.* 98, 10362.
- Moir, S., Malaspina, A., Pickeral, O.K., Donoghue, E.T., Vasquez, J., Miller, N.J., Krishnan, S.R., Planta, M.A., Turney, K.F., Justement, J.S., Kottlil, S., Dyubul, M., Mican, J.M., Kovacs, C., Chun, T.W., Birse, C.E., Fauci, A.S., 2004. Decreased survival of B cells of HIV-viremic patients mediated by altered expression of receptors of the TNF superfamily. *J. Exp. Med.* 200, 587.
- Morris, L., Binley, J.M., Clas, B.A., Bonhoeffer, S., Astill, T.P., Kost, R., Hurley, A., Cao, Y., Markowitz, M., Ho, D.D., Moore, J.P., 1998. HIV-1 antigen-specific and -nonspecific B cell responses are sensitive to combination antiretroviral therapy. *J. Exp. Med.* 188, 233.
- Mosier, D.E., Gulizia, R.J., Baird, S.M., Wilson, D.B., 1988. Transfer of a functional human immune system to mice with severe combined immunodeficiency. *Nature* 335, 256.
- Mosier, D.E., Gulizia, R.J., Baird, S.M., Wilson, D.B., Spector, D.H., Spector, S.A., 1991. Human immunodeficiency virus infection of human-PBL-Scid mice. *Science* 251, 791.
- Murphy, W.J., Taub, D.D., Longo, D.L., 1996. The huPBL-SCID mouse as a means to examine human immune function in vivo. *Semin. Immunol.* 8, 233.
- Ngo-Giang-Huong, N., Deveau, C., Da Silva, I., Pellegrin, I., Venet, A., Harzic, M., Venet, A., Harzic, M., Sinet, M., Delfraissy, J.F., Meyer, L., Goujard, C., Rouzioux, C., 2001. Proviral HIV-1 DNA in subjects followed since primary HIV-1 infection who suppress plasma viral load after one year of highly active antiretroviral therapy. *AIDS* 15, 665.
- Samuelsson, A., Sonnerborg, A., Heuts, N., Coster, J., Chiodi, F., 1997. Progressive B cell apoptosis and expression of Fas ligand during human immunodeficiency virus type 1 infection. *AIDS Res. Hum. Retrovir.* 13, 1031.
- Schmidt, G., Amiraian, K., Frey, H., Wethers, J., Stevens, R.W., Berns, D.S., 1989. Monitoring human immunodeficiency virus type 1-infected patients by ratio of antibodies to gp41 and p24. *J. Clin. Microbiol.* 27, 843.
- Schoonjans, F., Zalata, A., Depuydt, C.E., Comhaire, F.H., 1995. Medcalc: a new computer program for medical statistics. *Comput. Methods Program. Biomed.* 48, 257.
- Shirai, A., Cosentino, M., Leitman-Klinman, S.F., Klinman, D.M., 1992. Human immunodeficiency virus infection induces both polyclonal and virus-specific B cells activation. *J. Clin. Invest.* 89, 561.
- Shultz, L.D., Schweitzer, P.A., Christianson, S.W., Gott, B., Schweitzer, I.B., Tennent, B., McKenna, S., Mobraaten, L., Rajan, T.V., Greiner, D.L., Leiter, E.H., 1995. Multiple defects in innate and adaptive immunologic function in NOD/LtSz-scid mice. *J. Immunol.* 154, 180.
- Steyaert, S., Heyndrickx, L., Verhoye, L., Vermoesen, T., Donners, H., Franssen, K., Van Wanzele, F., Vandergucht, B., Vanham, G., Leroux-Roels, G., Vanlandschoot, P., in press. Inhibition of replication of primary HIV-1 isolates in huPBL-NOD/Scid mice by antibodies from HIV-1 infected patients. *Antiviral Res.*
- Strathdee, S.A., Frank, J.W., McLaughlin, J., Leblanc, M., Major, C., O'Shaughnessy, M.V., Read, S.E., 1995. Quantitative measures of human immunodeficiency virus-specific antibodies predict progression to AIDS. *J. Infect. Dis.* 172, 1375.
- Tanaka, T., Tsudo, M., Karasuyama, H., Kitamura, F., Kono, T., Hatakeyama, M., Taniguchi, T., Miyasaka, M., 1991. A novel monoclonal antibody against murine IL-2 receptor beta-chain. Characterization of receptor expression in normal lymphoid cells and EL-4 cells. *J. Immunol.* 147, 2222.
- Tanaka, T., Kitamura, F., Nagasaka, Y., Kuida, K., Suwa, H., Miyasaka, M., 1993. Selective long-term elimination of natural killer cells in vivo by an anti-interleukin 2 receptor beta chain monoclonal antibody in mice. *J. Exp. Med.* 178, 1103.
- Titanji, K., De Milito, A., Cagigi, A., Thorstensson, R., Grützmeier, S., Atlas, A., Hejdeman, B., Kroon, F.P., Lopalco, L., Nilsson, A., Chiodi, F., 2006. Loss of memory B cells impairs maintenance of long-term serologic memory during HIV-1 infection. *Blood* 108, 1580.
- Tournoy, K.G., Depraetere, S., Pauwels, R.A., Leroux-Roels, G.G., 2000. Mouse strain and conditioning regimen determine survival and function of human leucocytes in immunodeficient mice. *Clin. Exp. Immunol.* 119, 231.
- Traggiai, E., Becker, S., Subbarao, K., Kolesnikova, L., Uematsu, Y., Gismondo, M.R., Murphy, B.R., Rappuoli, R., Lanzavecchia, A., 2004. An efficient method to make human monoclonal antibodies from memory B cells: potent neutralization of SARS coronavirus. *Nat. Med.* 10, 871.
- Trkola, A., Kuster, H., Leemann, C., Oxenius, A., Fagard, C., Furrer, H., Battegay, M., Vernazza, P., Bernasconi, E., Weber, R., Hirschel, B., Bonhoeffer, S., Gunthard, H.F., 2004. Humoral immunity to HIV-1: kinetics of antibody responses in chronic infection reflects capacity of immune system to improve viral set point. *Blood* 104, 1784.
- Vanlandschoot, P., Van Houtte, F., Roobrouck, A., Farhoudi, A., Leroux-Roels, G., 2002. Hepatitis B surface antigen suppresses the activation of monocytes through interaction with a serum protein and a monocyte-specific receptor. *J. Gen. Virol.* 83, 1281.
- Yarchoan, R., Redfield, R.R., Broder, S., 1986. Mechanisms of B cell activation in patients with acquired immunodeficiency syndrome and related disorders. *J. Clin. Invest.* 78, 439.
- Zwart, G., van der Hoek, L., Valk, M., Cornelissen, M.T., Baan, E., Dekker, J., Koot, M., Kuiken, C.L., Goudsmit, J., 1994. Antibody responses to HIV-1 envelope and gag epitopes in HIV-1 seroconverters with rapid versus slow disease progression. *Virology* 201, 285.