

REVIEW

Open Access

Persistent digestive disorders in the tropics: causative infectious pathogens and reference diagnostic tests

Sören L Becker^{1,2,3}, Jürg Vogt^{1,2}, Stefanie Knopp^{1,2}, Marcus Panning⁴, David C Warhurst⁵, Katja Polman⁶, Hanspeter Marti^{2,7}, Lutz von Müller³, Cedric P Yansouni^{8,9}, Jan Jacobs⁹, Emmanuel Bottieau⁹, Moussa Sacko¹⁰, Suman Rijal¹¹, Fransiska Meyanti¹², Michael A Miles⁵, Marleen Boelaert¹³, Pascal Lutumba^{14,15}, Lisette van Lieshout¹⁶, Eliézer K N'Goran^{17,18}, François Chappuis¹⁹ and Jürg Utzinger^{1,2*}

Abstract

Background: Persistent digestive disorders account for considerable disease burden in the tropics. Despite advances in understanding acute gastrointestinal infections, important issues concerning epidemiology, diagnosis, treatment and control of most persistent digestive symptomatologies remain to be elucidated. Helminths and intestinal protozoa are considered to play major roles, but the full extent of the aetiologic spectrum is still unclear. We provide an overview of pathogens causing digestive disorders in the tropics and evaluate available reference tests.

Methods: We searched the literature to identify pathogens that might give rise to persistent diarrhoea, chronic abdominal pain and/or blood in the stool. We reviewed existing laboratory diagnostic methods for each pathogen and stratified them by (i) microscopy; (ii) culture techniques; (iii) immunological tests; and (iv) molecular methods. Pathogen-specific reference tests providing highest diagnostic accuracy are described in greater detail.

Results: Over 30 pathogens may cause persistent digestive disorders. Bacteria, viruses and parasites are important aetiologic agents of acute and long-lasting symptomatologies. An integrated approach, consisting of stool culture, microscopy and/or specific immunological techniques for toxin, antigen and antibody detection, is required for accurate diagnosis of bacteria and parasites. Molecular techniques are essential for sensitive diagnosis of many viruses, bacteria and intestinal protozoa, and are increasingly utilised as adjuncts for helminth identification.

Conclusions: Diagnosis of the broad spectrum of intestinal pathogens is often cumbersome. There is a need for rapid diagnostic tests that are simple and affordable for resource-constrained settings, so that the management of patients suffering from persistent digestive disorders can be improved.

Keywords: Bacteria, Clinical microbiology, Diagnosis, Digestive disorders, Gastroenterology, Helminths, Intestinal protozoa, Persistent diarrhoea, Virus

Background

Diarrhoeal diseases and other digestive disorders are leading causes of morbidity and mortality worldwide, with the highest burden concentrated in tropical and subtropical areas that often lack access to clean water and adequate sanitation, and where hygienic conditions

are generally poor [1]. According to the World Health Organization (WHO), diarrhoea is classified into three different categories, namely (i) acute watery diarrhoea (lasting several hours or days); (ii) acute bloody diarrhoea (synonymous: dysentery); and (iii) persistent diarrhoea (lasting 14 days or longer) [2]. 'Chronic diarrhoea' is often referred to as an individual term applicable to diarrhoea lasting more than 4–6 weeks, but it still lacks an unambiguous definition.

With an estimated burden of 89.5 million disability-adjusted life years (DALYs) caused in 2010, diarrhoeal

* Correspondence: juerg.utzinger@unibas.ch

¹Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel, Switzerland

²University of Basel, Basel, Switzerland

Full list of author information is available at the end of the article

diseases rank fourth in the recently published Global Burden of Disease Study [3]. Acute diarrhoeal episodes are mainly due to bacterial and viral pathogens that may cause a variety of clinical syndromes ranging from self-limiting events to life-threatening diseases. Children are most vulnerable and diarrhoeal diseases were responsible for more than 1.4 million deaths in 2010, ranking this disorder at position seven on the main causes of death [4]. In the last decades, concerted efforts have considerably improved our understanding of the epidemiology, diagnosis, treatment and control of many diarrhoeagenic pathogens globally, for instance due to the introduction of rotavirus vaccination programmes in many countries since 2006 [5]. As a result, mortality due to diarrhoeal diseases has been reduced from an estimated 2.5 million in 1990 to just under 1.5 million in 2010, a decrease of 42% [4]. However, few research activities have focused on the investigation of persistent diarrhoea and non-acute abdominal pain due to digestive disorders in the tropics. Hence, little is known regarding its aetiology, epidemiology and disease burden. It is widely acknowledged that intestinal parasites, particularly helminths and intestinal protozoa play a major role as causative agents of persistent digestive symptomatology [6].

Infections with helminths and intestinal protozoa belong to the neglected tropical diseases, along with other diseases caused by bacterial (e.g. Buruli ulcer), viral (e.g. dengue) and fungal infections (e.g. mycetoma) [7]. More than 5 billion people are at risk of neglected tropical diseases, with the common soil-transmitted helminths (i.e. *Ascaris lumbricoides*, hookworm and *Trichuris trichiura*), exhibiting the widest geographical distribution [8]. Due to their intimate connection with poverty, the highest prevalences of neglected tropical diseases are observed in remote rural and deprived urban settings in the developing world [7,9,10]. Neglected tropical diseases drain the social and economic development in endemic countries and they negatively impact on people's quality of life and well-being at all levels [11-15].

A major challenge in the clinical management of persistent digestive disorders is the weakness of health systems in many low-income countries [16-18]. Hence, affected people might only seek care at a late stage in their therapeutic itinerary, usually at primary health care centres [19,20]. However, these primary health care centres are often under-staffed and ill-equipped, resulting in a low quality of care. The causes of persistent diarrhoea and other digestive disorders are frequently misdiagnosed due to the often unspecific clinical presentations and the absence of evidence-based algorithms for in-depth investigation [7,21]. The notorious underfinancing of health systems in many tropical and subtropical countries explains the severe neglect of laboratory networks and the only limited availability of basic tests for diagnostic

services (e.g. direct faecal smears for helminth diagnosis or blood films for malaria diagnosis) [22]. Hence, in many developing countries, current diagnostic and treatment algorithms are often empirical, whereas local prevalence data and differential diagnoses are rarely taken into account at the primary care level.

Against this background, NIDIAG, an international collaboration on integrated diagnosis-treatment platforms, funded by the European Commission, sets out to develop an improved system for delivering primary health care in resource-constrained settings and proposes an integrated approach to this challenge. Emphasis is placed on a patient-centred approach starting from the presentation at the primary health care level of a clinical syndrome that might be due to 'common' pathogens. Three clinical syndromes will be investigated in the NIDIAG framework, namely (1) neurological disorders [23]; (2) persistent fever [24]; and (3) digestive disorders. Here, we focus on digestive disorders, which we define as (i) persistent (≥ 2 weeks) abdominal pain; (ii) persistent (≥ 2 weeks) diarrhoea (dysenteric or non-dysenteric); and/or (iii) blood in the stool. These digestive disorders will be investigated at different study sites in tropical areas of Africa (Côte d'Ivoire and Mali) and Asia (Indonesia and Nepal). Before clinically applicable diagnosis-treatment algorithms can be developed, the following major challenges/open issues have to be addressed. Firstly, few studies analysed the spectrum of intestinal pathogens causing persistent digestive disorders in the tropics. Therefore, epidemiological investigations targeting all potential pathogens are desirable to define the most common bacteria, parasites and viruses in the different study settings. Secondly, most diagnostic tests have only been validated in Western settings, and hence their diagnostic accuracy in the tropics remains to be determined.

In this manuscript, pursuing an extensive literature review complemented with expert opinion, we provide an overview of potential pathogens (bacterial, parasitic and viral) that might give rise to digestive disorders as defined above. Available diagnostic tests for the identified pathogens are summarised and reviewed, and we propose pathogen-specific reference tests to be utilised for an in-depth diagnostic work-up of symptomatic patients in the different study sites.

Methods

Framework

A symptomatology according to the aforementioned inclusion criteria for the syndrome of digestive disorders is likely to be caused by a large variety of infections, but also non-infectious diseases. For example, blood in the stool, accompanied by persistent abdominal pain, may be indicative of colorectal carcinoma or inflammatory disorders (e.g. Crohn's disease and ulcerative colitis), but

may also be a sign of *Schistosoma mansoni* (a helminth) or *Entamoeba histolytica* (an intestinal protozoon) infection. The aim of the NIDIAG project is to develop evidence-based diagnosis-treatment algorithms that can easily be applied in resource-constrained health care settings. As neither diagnosis nor treatment of many non-infectious diseases are currently feasible in remote rural areas of most developing countries, only infectious aetiologies of digestive disorders that may cause severe disease and that are treatable will be thoroughly assessed within the frame of the NIDIAG project and were therefore prioritised in our literature search.

Search strategy, data extraction and analysis

We performed a literature review to identify and define the bacterial, parasitic and viral pathogens that may give rise to persistent diarrhoea and chronic digestive disorders, and to obtain information on their respective diagnostic methods in order to describe appropriate reference laboratory tests. Since the role of fungi as causative pathogens of gastrointestinal infections is still under debate, fungal infections were not included in this review [25]. The available literature was reviewed by three independent groups. The results were compared, discussed and finally synthesised. Additionally, a number of experts were consulted to complement the literature review.

In a first step, we examined a series of textbooks pertaining to medical bacteriology, parasitology and virology. Moreover, we searched the electronic database MEDLINE/PubMed for infectious pathogens that may cause digestive disorders as defined in the inclusion criteria. After having identified a set of more than 30 pathogens, we searched the database with the following search term for all infectious agents: “disease name/[Mesh]” and the subheading “diagnosis” (e.g. “ascariasis/diagnosis” [Mesh]). The focus of the MEDLINE/PubMed search was on established laboratory tests as well as on newer diagnostic methods, which have been validated recently or are currently under validation (e.g. studies objectively assessing the diagnostic accuracy of different tests). Hence, we primarily focused on reviews, comparative studies and evaluation studies. Our search had no language or other restrictions and we included studies that were published until mid-April 2012.

Results

Our literature review revealed more than 30 bacterial, parasitic and viral pathogens that may cause persistent digestive disorders. Many of these infectious agents are epidemiologically well characterised in Western settings, while data regarding their occurrence in tropical and subtropical areas are scarce and often contradictory [6,26-28]. Table 1 provides a list of all selected pathogens and typical clinical characteristics that may assist clinicians to curtail

their differential diagnosis. However, pathogen-specific diagnosis can rarely be done based on the clinical presentations, and hence additional diagnostic tools are needed.

The large number of available diagnostic tests for the selected pathogens is a challenge for providing the single most accurate method for a given pathogen. Hence, we classified the different methods into four diagnostic categories, namely (i) microscopy; (ii) culture; (iii) immunology (including enzyme immunoassays (EIA), serotyping of isolates and serology); and (iv) molecular biological diagnosis (e.g. polymerase chain reaction (PCR) assays and DNA sequencing). Selection of a reference test for each specific pathogen is primarily based on the sensitivity and specificity of the test as well as practical considerations (e.g. costs, ease of application, availability, etc.). The results are presented in Table 2 (bacteria), Table 3 (intestinal protozoa), Table 4 (helminths) and Table 5 (viruses). Specific issues on the diagnostic work-up are summarised in the following sub-chapters.

Bacterial pathogens

Aeromonas spp., *Campylobacter jejuni*, *C. coli*, *Plesiomonas shigelloides*, *Salmonella enterica* (non-typhoidal serovars, e.g. *S. enterica* ser. *Enteritidis*, *S. enterica* ser. *Typhimurium*), *Shigella* spp., *Vibrio* spp., *Yersinia enterocolitica*, *Y. pseudotuberculosis*

A stool culture on selective media is the test of choice to detect these diarrhoeagenic bacteria [31,48,50,52,53]. Different solid media (e.g. selective agar plates containing antibiotics and substances favouring the growth of the sought microorganism) are inoculated with a stool specimen to detect and isolate enteric pathogens. The additional use of a selective enrichment broth is helpful to identify pathogenic bacteria if their presence is quantitatively so low that they might otherwise be overlooked on solid media due to the overgrowth of non-pathogenic intestinal flora. The inoculated media are usually incubated for 24–72 hours at 35°C at ambient atmosphere to allow the bacteria to form macroscopically visible colonies. Of note, *Campylobacter* spp. are isolated using different growth conditions, i.e. incubation at a higher temperature of 42°C in microaerophilic atmosphere [53].

Following the incubation period, the agar plates are examined and morphologically suspicious colonies are identified using different biochemical identification panels or automated phenotypic identification systems (e.g. Vitek[®]; bioMérieux, Marcy l'Étoile, France). Recently, more rapid identification algorithms making use of mass spectrometry (MS) have successfully been implemented into clinical microbiology laboratories (e.g. matrix-assisted laser desorption ionization time-of-flight (MALDI-TOF) MS (e.g. MicroFlex LT; Bruker Daltonics, Bremen, Germany) [98].

Stool culture remains the diagnostic ‘gold’ standard for enteric pathogenic bacteria disposing certain characteristics

Table 1 Overview of intestinal pathogens (bacteria, intestinal protozoa, helminths and viruses) that may cause persistent digestive disorders in infected individuals

Enteric pathogen	Persistent diarrhoea	Persistent abdominal pain	Blood in the stool
Bacteria			
<i>Aeromonas</i> spp.	+	-	-
<i>Campylobacter jejuni</i> , <i>C. coli</i>	+	+	+
<i>Clostridium difficile</i>	+	+	+
<i>Escherichia coli</i>			
Enteroaggregative <i>E. coli</i> (EAEC)	+	+	+
Enteropathogenic <i>E. coli</i> (EPEC)	+	+	-
Enteroinvasive <i>E. coli</i> (EIEC)	+	+	+
Enterohaemorrhagic <i>E. coli</i> (STEC/EHEC)	+	+	+
Enterotoxigenic <i>E. coli</i> (ETEC)	+	+	-
Diffusely adherent <i>E. coli</i> (DAEC)	+	-	-
<i>Mycobacterium tuberculosis</i> and atypical mycobacteria	+	+	+
<i>Plesiomonas shigelloides</i>	+	-	-
<i>Salmonella enterica</i> (typhoidal and non-typhoidal serovars)	+	+	+
<i>Shigella</i> spp.	+	+	+
<i>Tropheryma whipplei</i>	+	-	-
<i>Vibrio</i> spp.	+	-	-
<i>Yersinia enterocolitica</i> , <i>Y. pseudotuberculosis</i>	+	-	-
Intestinal protozoa			
<i>Balantidium coli</i>	+	+	+
<i>Blastocystis hominis</i> ^a	(+)	(+)	-
<i>Cryptosporidium</i> spp.	+	+	-
<i>Cyclospora cayetanensis</i>	+	-	-
<i>Dientamoeba fragilis</i> ^a	+	+	-
<i>Entamoeba histolytica</i>	+	+	+
<i>Giardia intestinalis</i> (syn.: <i>G. lamblia</i> and <i>G. duodenalis</i>)	+	+	-
<i>Isospora belli</i> (syn.: <i>Cystoisospora belli</i>)	+	(+)	-
Species of microsporidia	+	+	-
Helminths			
Cestodes			
<i>Diphyllobothrium latum</i>	+	+	-
<i>Hymenolepis</i> spp.	+	-	-
<i>Taenia</i> spp.	+	+	-
Nematodes			
<i>Ascaris lumbricoides</i>	+	+	-

Table 1 Overview of intestinal pathogens (bacteria, intestinal protozoa, helminths and viruses) that may cause persistent digestive disorders in infected individuals (Continued)

<i>Capillaria philippinensis</i>	+	+	-
Hookworm (<i>Ancylostoma duodenale</i> and <i>Necator americanus</i>)	+	+	-
<i>Strongyloides stercoralis</i>	+	+	(+)
<i>Trichuris trichiura</i>	+	+	-
Trematodes			
Intestinal flukes	+	+	-
Intestinal blood flukes: <i>Schistosoma mansoni</i> , <i>S. intercalatum</i> , <i>S. japonicum</i> , <i>S. mekongi</i>	+	+	+
Viruses			
Adenovirus	+	-	-
Astrovirus	(+)	-	-
Bocavirus	(+)	-	-
Coronavirus	(+)	-	-
Cytomegalovirus (CMV)	+	+	+
Enterovirus	+	-	-
Human immunodeficiency virus (HIV-1/2)	+	+	-
Norovirus	+	-	-
Parechovirus	(+)	-	-
Rotavirus	+	+	-
Sapovirus	(+)	-	-

+, existing risk; (+), low risk; -, no risk.

^a There is an ongoing debate whether these intestinal protozoa have pathogenic potential or should rather be seen as simple commensals of the gastrointestinal tract [29,30].

which enable them to be selected out of the normal gastrointestinal flora, while other bacteria without such characteristics cannot be distinguished from apathogenic gut bacteria by culture methods alone (see below). Stool culture has important advantages, such as testing of isolated pathogens for antimicrobial susceptibility. As the successful antibiotic treatment of many bacterial intestinal infections requires knowledge of local resistance patterns (e.g. extent of fluoroquinolone-resistant *Campylobacter* strains), stool culture techniques remain mandatory to guide therapeutic interventions. However, these tests are laborious and require experienced personnel, and typically take 48–72 hours to obtain first results. Hence, other, more rapid diagnostic tests (RDTs) have been developed for some pathogens. For *Campylobacter* spp., for example, EIAs detecting a specific antigen in the stool proved to be a sensitive alternative to stool culture with results available within a few hours [32,99]. However, there is no international consensus on

immunological assays for detection of *Campylobacter* spp. and no globally validated and standardised approach, so that these tools should not replace the selective stool culture [100]. PCR assays, characterised by high sensitivity and specificity, have been developed for most of the aforementioned bacteria. Thus far, however, integration into clinical routine testing is still limited. Important drawbacks are the high cost, the need for sophisticated laboratory equipment and well-trained laboratory technicians. Moreover, PCR cannot distinguish between dead or alive bacteria and does not allow testing for antimicrobial susceptibility. However, newly developed multiplex PCR assays are increasingly being evaluated as fast screening tests for early detection of various important enteric pathogens. Besides PCR, novel molecular diagnostics are currently being developed and validated for many bacterial and viral pathogens, e.g. loop-mediated isothermal amplification (LAMP) assays. Results obtained thus far are promising [101,102], but it remains to be elucidated whether such nucleic acid amplification tests can be employed on a larger scale in resource-constrained settings in the tropics.

Salmonella enterica ser. Typhi/Paratyphi

Diagnosis of enteric fever is challenging and often delayed or not performed due to the unavailability of the most sensitive techniques in areas of high endemicity [103]. In these settings, the Widal test (measuring an increasing *S. Typhi*-specific antibody titer over the course of 10 days in patient serum samples) is often the only available test, despite its poor diagnostic performance. Contrary to non-typhoidal salmonellosis, stool culture is not sufficiently sensitive to diagnose infection due to *S. Typhi/S. Paratyphi*. Culturing blood and bone marrow is more sensitive, but bone marrow aspiration is only rarely performed in tropical areas due to a lack of adequately equipped hospitals and laboratories [104]. Blood cultures should be obtained during the first week of disease to achieve adequate sensitivity [45]. Serotyping of isolates (e.g. by agglutination of Vi antigen or rapid detection of various antigens or IgM antibodies by different EIA kits) is helpful for a timely diagnosis, but lacks sensitivity and specificity [46]. PCR assays have been developed for different antigens of invasive *S. enterica* serovars, but still need further development and validation before they can be more widely recommended [105].

Clostridium difficile

C. difficile can be found as part of the physiological intestinal flora, but toxin-producing strains may cause severe diarrhoea, which is most frequently seen in hospitalised patients who recently received antibiotic treatment [33]. A selective stool culture (toxigenic culture, performed on a selective agar medium or after ethanol shock pretreatment) followed by tests for toxin production

remains the diagnostic 'gold' standard for *C. difficile* [33] and is particularly useful when the quantity of toxins in stool samples is small [34]. A laborious and technically difficult cell culture cytotoxicity assay is still regarded as an alternative reference standard, but is seldom performed in most microbiological laboratories. More recently developed PCR assays targeting a toxin-encoding gene are currently discussed as an alternative method for early diagnosis of *C. difficile* infection. Such molecular methods allow a more precise characterisation of isolated *C. difficile* strains, e.g. ribotype differentiation [35,106,107]. Sensitivity and specificity of PCR have been reported to vary between 85% and 100% [108]. However, various molecular assays exist which are not yet fully standardised, and the diagnostic performance of commercially available kits may differ considerably from in-house molecular testing methods in use at different laboratories. Of note, PCR can only prove the presence of the toxin-encoding gene, but cannot distinguish between asymptomatic carriage and acute infection.

In clinical practice, an easily applicable two-step approach is recommended for rapid and reasonably sensitive diagnosis of *C. difficile* [109]. Firstly, a screening test for *C. difficile*-associated glutamate dehydrogenase (GDH) should be performed to indicate the bacterium's presence in a stool sample. If positive, it should be followed by a test for toxin production (e.g. toxin A/B EIA). This procedure does not require an extensively equipped laboratory and generates accurate results within a few hours. However, the sensitivity and specificity of this two-step approach are limited, and hence toxigenic culture and PCR testing should always be performed when there is a high clinical suspicion despite negative test results [110].

Pathogenic Escherichia coli strains

Diagnosis of pathogenic *E. coli* is challenging, as these bacteria constitute an important part of the physiological intestinal flora and only some strains have diarrhoeagenic potential [41]. There are at least six groups of pathogenic *E. coli* strains, namely (i) diffusely adherent (DAEC); (ii) enteroaggregative (EAEC); (iii) enterohaemorrhagic (EHEC, including STEC = shiga toxin-producing *E. coli*); (iv) enteroinvasive (EIEC); (v) enteropathogenic (EPEC); and (vi) enterotoxigenic *E. coli* (ETEC). Pathogenic *E. coli* strains that carry simultaneously virulence factors from different pathotypes may cause severe clinical outbreaks. In mid-2011 in Germany, for example, the *E. coli* strain O104:H4 (an EAEC capable of EHEC/STEC-specific shiga toxin production) caused 2,987 cases of acute, often severe gastroenteritis and 855 cases of haemolytic-uraemic syndrome which led to 53 deaths [111].

While diagnostic procedures are poorly standardised for the pathotypes DAEC and EAEC, molecular

Table 2 Diagnostic tests for important bacterial pathogens that may cause persistent digestive disorders

Infectious pathogen	Diagnostic method				
	Microscopy	Stool culture	Immunology	Molecular biology (PCR)	Reference(s)
<i>Aeromonas</i> spp.	- ^a	Culture on cefsulodin-irgasan-novobiocin (CIN) or selective <i>Aeromonas</i> agar	-	(Experimental, not validated)	[31]
<i>Campylobacter jejuni</i> , <i>C. coli</i>	Darkfield microscopy: motile, curved or S-shaped rods (suggestive of <i>Campylobacter</i> spp.)	Culture on selective medium^b (42°C, microaerophilic conditions)	<ul style="list-style-type: none"> • Faecal antigen enzyme immunoassay: <i>Campylobacter</i>-specific antigen (SA) • Serology (important for diagnosis of postinfectious immunological diseases) 	<i>hipO</i> gene (<i>C. jejuni</i>), <i>glyA</i> gene (<i>C. coli</i>)	[32]
<i>Clostridium difficile</i>	- ^c	Culture on selective medium, e.g. cycloserin-cefoxitin-fructose agar (CCF) + toxigenic culture	<ul style="list-style-type: none"> • 2-step algorithm: 1) Screening: EIA for glutamate dehydrogenase (GDH) 2) ELISA for detection of toxin A and B • Cell cytotoxicity assay for detection of toxin A and B 	Toxin genes (increasingly being used in clinical routine)	[33-35]
<i>Escherichia coli</i> Enteroaggregative <i>E. coli</i> (EAEC)	- ^a	HEp-2 cell adherence assay (following incubation in Luria broth)	<ul style="list-style-type: none"> • Serology: antibody response against Plasmid-encoded toxin (<i>Pet</i>) • ELISA: secretory immunoglobulin A response to EAEC 	<i>AggR</i> , <i>CVD432</i> , <i>EAST1</i> (most common virulence factors, not always present)	[36]
Enteropathogenic <i>E. coli</i> (EPEC)	- ^a	Culture on MacConkey (MAC) agar	-	<i>eae</i> gene	[37]
Enteroinvasive <i>E. coli</i> (EIEC)	- ^a	Culture on MAC agar	ELISA: detection of the <i>ipaC</i> gene	<i>ipaH</i>, <i>ipaB</i> genes	[38]
Enterohaemorrhagic <i>E. coli</i> (EHEC including STEC)	- ^a	Culture on sorbitol-MAC agar (most O157: H7 strains form sorbitol-negative colonies)	<ul style="list-style-type: none"> • O157 latex agglutination test • Shiga toxins 1 & 2 (ELISA) 	STEC: <i>stx1</i>, <i>stx2</i> genes EHEC: <i>stx1/stx2</i> + <i>eae</i> gene	[39,40]
Enterotoxigenic <i>E. coli</i> (ETEC)	- ^a	Culture on MAC agar	Several immunoassays for toxin detection	<i>stla/stlb</i> and <i>lt</i> genes	[41]
Diffusely adherent <i>E. coli</i> (DAEC)	- ^a	HEp-2 cell adherence assay (following incubation in Luria broth)	-	<i>daaD</i> gene	[42]
<i>Mycobacterium tuberculosis</i> and atypical mycobacteria	- Histopathological examination of intestinal biopsies - Acid-fast stain (e.g. Ziehl-Neelsen, Kinyoun, Auramin)	Culture of biopsy material	<ul style="list-style-type: none"> • Interferon-gamma-release assay (IGRA) on heparinised blood samples • Tuberculin skin test 	Nucleic acid amplification tests (lacks sensitivity for diagnosis of extrapulmonary tuberculosis)	[43,44]
<i>Plesiomonas shigelloides</i>	- ^a	Culture on CIN agar	-	-	

Table 2 Diagnostic tests for important bacterial pathogens that may cause persistent digestive disorders (Continued)

Infectious pathogen	Diagnostic method				Reference(s)
	Microscopy	Stool culture	Immunology	Molecular biology (PCR)	
<i>Salmonella enterica</i> (typhoidal and non-typhoidal serovars)	- ^a	<ul style="list-style-type: none"> • Culture^d from blood and/or bone marrow (enteric fever) • Culture^d from stool or duodenal aspirate (typhoidal and non-typhoidal salmonellosis) 	<ul style="list-style-type: none"> • Serotyping of isolates (Vi antigen) • ELISA: detection of <i>S. typhi</i> antigens (blood) • Widal agglutination test (commonly used in Africa) 	(Mainly for research purpose)	[45-47]
<i>Shigella dysenteriae</i> , <i>S. flexneri</i> , <i>S. boydii</i> , <i>S. sonnei</i>	- ^a	Culture on MAC, XLD, HE or Leifson agar	Agglutination tests to detect serogroup and serotype	<i>ipaH</i> , <i>ipl</i> genes	[48]
<i>Tropheryma whipplei</i>	Histopathological examination of PAS-stained intestinal biopsies: sickleform particle-containing cells	(Only in highly specialised laboratories)	Immunohistochemistry on PAS-positive biopsy material	<i>whip1</i>, <i>whip2</i> genes	[49]
<i>Vibrio</i> spp.	Darkfield microscopy: comma-shaped, motile bacteria (highly suggestive of <i>Vibrio</i> spp.)	Culture on TCBS agar	-	PCR for species differentiation (<i>V. cholerae</i> , <i>V. parahaemolyticus</i> , <i>V. vulnificus</i>)	[50,51]
<i>Yersinia enterocolitica</i> , <i>Y. pseudotuberculosis</i>	- ^a	Culture on CIN agar	Serology (important for diagnosis of postinfectious immunological diseases)	PCR (reference laboratories and research purposes)	[52]

The laboratory techniques are divided into different categories and recommended tests for each pathogen are highlighted.

^a Gram staining of stool samples can be useful to evaluate the presence of leucocytes, but is not helpful to differentiate between pathogenic bacteria and apathogenic microbial flora.

^b Commonly employed selective media for detection of *Campylobacter* spp. include charcoal-cefoperazone-deoxycholate agar, *Campylobacter* blood agar plate, and cefoperazone-vancomycin-amphotericin agar [53].

^c Detection of *C. difficile* in the Gram stain is not adequate to differentiate between clinical infection and simple colonisation with *C. difficile* [54].

^d Commonly employed selective media for growth of *S. enterica* are MAC, XLD, HE, Leifson agar or other chromogenic media.

biological testing has revolutionized the diagnostic algorithms for the other diarrhoeagenic *E. coli*. Modern multiplex PCR assays targeting unique genes of EHEC/STEC, EIEC, EPEC and ETEC allow a rapid molecular characterisation of these pathogenic strains. Hence, multiplex PCR assays have become the test of choice with excellent sensitivity and specificity (>99%) [42]. Indeed, these tests have overcome important drawbacks of the classical stool culture, which often detects only some important strains (e.g. in the case of EHEC the O157:H7 strain on Sorbitol-MacConkey agar), but misses others that lack characteristic biochemical properties [39]. However, the integration of such multiplex PCR assays into routine testing of clinical samples remains restricted to well-equipped laboratories, and hence, these molecular techniques are only rarely available in endemic settings in the tropics.

***Mycobacterium tuberculosis* and atypical mycobacteria (e.g. *M. avium*)**

Gastrointestinal tuberculosis is the sixth most common manifestation of extrapulmonary tuberculosis and causes

considerable morbidity, including persistent diarrhoea and abdominal pain [112]. Atypical mycobacteria (synonymous: mycobacteria other than tuberculosis, MOTT), particularly *M. avium*, are an important cause of long-lasting diarrhoea and gastrointestinal complaints in HIV-infected individuals. Accurate diagnosis is difficult and relies on in-depth analysis of intestinal biopsy specimens by histopathological examination, microscopy after acid-fast staining (e.g. Ziehl-Neelsen, Auramin or Kinyoun techniques) and culture on selective media suitable for mycobacteria. Unless performed using oil immersion, histopathology often fails to distinguish between gastrointestinal tuberculosis and other granulomatous disorders, such as Crohn's disease [113,114]. An important drawback when culturing mycobacteria is their slow growth; it might take up to six weeks until cultures become positive. However, culture is the most sensitive technique and remains the diagnostic 'gold' standard [115]. Different molecular biological assays have been developed for various mycobacteria, but lack sensitivity for extrapulmonary tuberculosis and have not yet been validated for gastrointestinal tuberculosis [43].

Table 3 Diagnostic tests for important intestinal protozoa that may cause persistent digestive disorders

Infectious pathogen	Diagnostic method				Reference(s)
	Microscopy	Stool culture	Immunology	Molecular biology (PCR)	
<i>Balantidium coli</i>	Stool microscopy <ul style="list-style-type: none"> • Wet mount smears (unstained or iodine stain) • Concentration techniques (e.g. formalin-ether) • Permanent stains (e.g. with iron hematoxylin) 	-	-	-	[55]
<i>Blastocystis hominis</i>	Stool microscopy <ul style="list-style-type: none"> • Wet mount smears (unstained or iodine stain) • Permanent stains (e.g. with trichrome, iron hematoxylin, Giemsa) 	Stool culture on selective liquid media (no routine procedure, but beneficial in microscopically uncertain cases)	(No routine procedure)	(PCR mainly applied in research settings)	[56-58]
<i>Cryptosporidium</i> spp.	Stool microscopy <ul style="list-style-type: none"> • Wet mount smears (unstained or iodine stain) • Various staining techniques, especially acid-fast stains (e.g. Kinyoun, modified Ziehl-Neelsen) 	(No routine procedure)	<ul style="list-style-type: none"> • ELISA: faecal antigen detection • Fluorescence microscopy 	PCR (in reference laboratories and for species differentiation)	[59-62]
<i>Cyclospora cayentanensis</i>	Stool microscopy <ul style="list-style-type: none"> • Wet mount smears (light or epifluorescence microscopy) • Concentration techniques (e.g. formalin-ether) • Acid-fast stains (oocysts are variably acid-fast) 	(No routine procedure)	-	PCR (in reference laboratories)	[63]
<i>Dientamoeba fragilis</i>	Stool microscopy on stained smears (e.g. iron-hematoxylin, chlorazol black dye stain)	(No routine procedure)	-	PCR (in reference laboratories) on unpreserved stool samples	[64,65]
<i>Entamoeba histolytica</i>	Stool microscopy <ul style="list-style-type: none"> • Wet mount smears (trophozoites) • Formalin-ether concentration (cysts) • Permanent stains 	(No routine procedure)	<ul style="list-style-type: none"> • ELISA: faecal antigen detection able to distinguish between <i>E. histolytica</i> and <i>E. dispar/moshkovskii</i> (stool) • Serological antibody detection tests (blood samples) 	PCR (in reference laboratories)	[66-70]
<i>Giardia intestinalis</i> (syn.: <i>G. lamblia</i> and <i>G. duodenalis</i>)	Stool microscopy <ul style="list-style-type: none"> • Wet mount smears (trophozoites) • Formalin-ether concentration (cysts) • Permanent stains 	(No routine procedure)	<ul style="list-style-type: none"> • ELISA: faecal antigen detection 	PCR (in reference laboratories)	[60,71]

Table 3 Diagnostic tests for important intestinal protozoa that may cause persistent digestive disorders (Continued)

Infectious pathogen	Diagnostic method				Reference(s)
	Microscopy	Stool culture	Immunology	Molecular biology (PCR)	
<i>Isospora belli</i> (syn.: <i>Cystoisospora belli</i>)	Stool microscopy • Wet mount smears • Concentration techniques (e.g. formalin-ether) • Acid-fast stains	-	-	PCR (in reference laboratories)	[60,72,73]
Species of microsporidia (<i>Enterocytozoon bieneusi</i> , <i>Encephalitozoon</i> spp.)	• Transmission electron microscopy (gold standard, but not feasible as routine test) - Light microscopy (e.g. Uvitex B, Chromotrope R or Calcofluor White stain)	-	Serology: anti-microsporidial antibodies (indirect immunofluorescence assay)	PCR (in reference laboratories)	[60,74-76]

The laboratory techniques are divided into different categories and recommended tests for each pathogen are highlighted.

Tropheryma whipplei

Whipple's disease due to infection with *T. whipplei* is a rare disease characterised by chronic diarrhoea, wasting, abdominal pain, arthralgia and various other symptoms associated with organ involvement (e.g. encephalitis and endocarditis) [49]. The infectious agent was not identified until 1961 and many epidemiological and biological features still need to be elucidated [116]. Only highly specialised laboratories are able to grow *T. whipplei* on human fibroblast cells [117,118]. The development of a PCR assay targeting the genes *whip1* and *whip2* has been a major step forward and is nowadays the test of choice, especially in symptomatic patients without typical histopathological findings in intestinal biopsies (sickleform particle-containing cells on periodic acid-Schiff (PAS)-stained biopsy specimens) [49].

Parasitic pathogens: intestinal protozoa

***Balantidium coli*, *Blastocystis hominis*, *Cryptosporidium* spp., *Cyclospora cayetanensis*, *Dientamoeba fragilis*, *Entamoeba histolytica*, *Giardia intestinalis* (syn.: *G. lamblia* and *G. duodenalis*), *Isospora belli* (syn.: *Cystoisospora belli*), species of microsporidia**

The three main techniques for the diagnosis of human intestinal protozoan infections include (i) light microscopy; (ii) antigen detection (EIAs); and (iii) PCR assays. Since the first description of parasitic intestinal protozoa in human stools, documented by the Dutch microscopist Antony van Leeuwenhoek in 1681 [119], microscopic detection of protozoan cysts and trophozoites has been the most widely used diagnostic approach. On fresh stool samples, direct microscopy is performed by mixing a small amount of faeces with a physiological 0.9% sodium chloride (NaCl) solution. To increase sensitivity, various stool concentration

techniques have been developed, making use of either sedimentation or flotation with a formalin-ether concentration technique being the most widely used method in medical laboratories [120,121]. However, the formalin-ether concentration technique lacks sensitivity for several intestinal protozoan species as well as many helminths (described below), and hence there is a pressing need for new and more sensitive microscopic techniques (e.g. FLOTAC) [122] and non-microscopic diagnostics. Staining techniques can be helpful for microscopic parasite identification and might further improve the diagnostic accuracy. Indeed, some intestinal protozoan species require staining of the stool sample to be identified on microscopic examination. For example, acid-fast stains allow detection of *Cryptosporidium* spp., while species of microsporidia are best seen when using an Uvitex B or Calcofluor White stain. Still, correct identification of intestinal protozoan pathogens is challenging even for experienced laboratory technicians and for some species even impossible (e.g. *E. histolytica* based on cysts morphology). For *Cryptosporidium* spp., *E. histolytica* and *G. intestinalis*, sensitive EIAs detecting species-specific antigens in faecal samples have been developed, some of which are highly sensitive and complement microscopic stool examination in many clinical laboratories [123,124]. Especially for the diagnosis of *E. histolytica*, species differentiation based on alternative procedures is compulsory, since microscopy cannot readily distinguish between *E. histolytica* and the non-pathogenic *E. dispar* [66,125,126]. Of note, not all commercially available EIA antigen detection kits are *E. histolytica*-specific and some lack sensitivity, in particular if faecal samples have been stored for several days [67,127]. Over the past several years, highly sensitive PCR assays have been developed and standardised for many intestinal protozoan species.

Table 4 Diagnostic tests for important helminths that may cause persistent digestive disorders

Infectious pathogen	Diagnostic method				Reference(s)
	Microscopy	Stool culture	Immunology	Molecular biology (PCR)	
Cestodes					
<i>Diphyllobothrium latum</i>	Stool microscopy: identification of eggs or proglottids <ul style="list-style-type: none"> • Wet preparation • Ethyl-acetate or formalin-ether-based concentration techniques • Sedimentation techniques 	-	-	PCR and sequencing for species differentiation (for epidemiological purpose)	[77,78]
<i>Hymenolepis</i> spp.	Stool microscopy <ul style="list-style-type: none"> • Kato-Katz method • Ethyl-acetate or formalin-ether-based concentration techniques • Sedimentation techniques • FLOTAC 	-	-	PCR in research settings (for epidemiological purpose)	[79]
<i>Taenia</i> spp.	Stool microscopy <ul style="list-style-type: none"> • Perianal egg detection • (Graham's test applying adhesive tape) • Examination of tapeworms from purges 	-	<ul style="list-style-type: none"> • Coproantigen EIA • Serology: detection of specific circulating antibodies against <i>T. solium</i> 	PCR for species differentiation	[80]
Nematodes					
<i>Ascaris lumbricoides</i>	Stool microscopy: egg detection <ul style="list-style-type: none"> • Kato-Katz method • Ethyl-acetate or formalin-ether-based concentration techniques • Sedimentation techniques • FLOTAC 	-	-	PCR in research settings (for epidemiological purpose)	[81-83]
<i>Capillaria philippinensis</i>	Stool microscopy: egg detection <ul style="list-style-type: none"> • Ethyl-acetate or formalin-ether-based concentration techniques • Sedimentation techniques • (Kato-Katz method: great care is indicated to distinguish between <i>T. trichiura</i> and <i>C. philippinensis</i> eggs) 	-	-	-	[84,85]
Hookworms (<i>Ancylostoma duodenale</i> , <i>Necator americanus</i>)	Stool microscopy: egg detection <ul style="list-style-type: none"> • Kato-Katz method • Ethyl-acetate or formalin-ether-based concentration techniques • Sedimentation techniques • FLOTAC 	Culture on Koga agar and subsequent microscopic identification of larvae		PCR mainly applied in research settings (for epidemiological purpose)	[81-83]

Table 4 Diagnostic tests for important helminths that may cause persistent digestive disorders (Continued)

Infectious pathogen	Diagnostic method				
	Microscopy	Stool culture	Immunology	Molecular biology (PCR)	Reference(s)
<i>Strongyloides stercoralis</i>	<ul style="list-style-type: none"> • Stool: microscopy following Baermann funnel concentration • Microscopy of sputum, bronchoalveolar lavage, duodenal aspirate, skin biopsy 	Culture on Koga agar and subsequent microscopic identification of larvae	<ul style="list-style-type: none"> • ELISA tests detecting serum antibodies or faecal antigens • Indirect fluorescent antibody test 	PCR applied in research settings (for epidemiological purpose) and increasingly used for individual patient management	[86,87]
<i>Trichuris trichiura</i>	<ul style="list-style-type: none"> • Stool microscopy: egg detection • Kato-Katz method • Ethyl-acetate or formalin-ether-based concentration techniques • Sedimentation techniques • FLOTAC 	-	-	-	[81,82]
Trematodes					
Intestinal flukes	<ul style="list-style-type: none"> • Stool microscopy: egg detection • Kato-Katz method • Ethyl-acetate or formalin-ether-based concentration techniques • Stoll's dilution • Sedimentation techniques • FLOTAC 	-	ELISA to detect worm-specific antibodies or antigens in serum or stool	PCR applied in research settings (for epidemiological purpose)	[88]
Intestinal blood flukes: <i>Schistosoma mansoni</i> , <i>S. intercalatum</i> , <i>S. japonicum</i> , <i>S. mekongi</i>	<ul style="list-style-type: none"> • Stool microscopy: egg detection • Kato-Katz method • Ethyl-acetate or formalin-ether-based concentration techniques • Stoll's dilution • Sedimentation techniques • FLOTAC (first experiences for <i>S. mansoni</i>) • Miracidium-hatching test from stool samples 	-	<ul style="list-style-type: none"> - ELISA to detect serum antibodies or worm-specific antigens in serum or urine - RDT to detect CCA or CAA antigen in serum or urine (for <i>S. mansoni</i>) 	PCR applied in research settings for epidemiological purpose and increasingly used for individual patient management	[89]

The laboratory techniques are divided into different categories and recommended tests for each pathogen are highlighted.

Many of these assays (e.g. *Entamoeba* spp. differentiation by PCR) are currently being integrated into parasitological reference laboratories as an additional diagnostic tool to prove diagnosis in uncertain clinical cases [59,128,129]. Such molecular biological tools are of enormous importance to improve the correct species identification of many intestinal parasites, which are difficult to diagnose using conventional techniques [60,74].

Parasitic pathogens: helminths

Ascaris lumbricoides, *Capillaria philippinensis*, *Diphyllobothrium* spp., *Hymenolepis* spp., hookworm (*Ancylostoma duodenale* and *Necator americanus*), *Taenia* spp., *Trichuris trichiura*, intestinal flukes

Identification of helminth eggs on microscopic stool examination is the reference test for most intestinal helminth species. In hospitals and microbiological laboratories,

direct stool examination after prior concentration (e.g. by formalin-ether concentration technique) is most commonly employed, while the Kato-Katz thick smear technique is widely used in epidemiological studies and anthelmintic drug efficacy evaluations in endemic regions [81,130-132]. Direct microscopic examination is a cheap methodology, the microscope slides can rapidly be prepared for examination, and there is no need for sophisticated laboratory equipment. The eggs of most helminth species parasitising humans can easily be distinguished by a trained laboratory technician (see Figure 1 for eight selected helminth eggs). Hence, microscopy remains the standard reference test for *A. lumbricoides*, *T. trichiura*, hookworm, *Capillaria philippinensis*, *Diphyllobothrium* spp., *Hymenolepis* spp., *Taenia* spp. and blood flukes (*Schistosoma* spp.) [82,88,133,134]. However, microscopy is prone to a number of shortcomings. Firstly, microscopy is not very sensitive and especially infections of light intensity can be missed when only a single stool sample is analysed [131,135]. Multiple stool sampling, ideally over several consecutive days, increases the sensitivity [136], as well as the use of different concentration techniques, which are based on sedimentation (e.g. formalin-ether concentration technique), flotation or a combination of both (e.g. McMaster technique and FLOTAC) [135,137-139].

However, these techniques often require access to the power grid, a centrifuge and different chemical reagents, which are not always available in tropical settings. Moreover, the diagnostic sensitivity for different helminth species often varies considerably, and no currently available concentration technique is able to concurrently detect intestinal protozoa and helminths with the same diagnostic accuracy [122,140].

Secondly, microscopy results heavily depend on the quality of the slide preparation and on the experience of the laboratory technician reading the slides. Thirdly, the eggs of some helminth species such as the two hookworm species *A. duodenale* and *N. americanus* are virtually indistinguishable by microscopy. Fourthly, the nematode *Strongyloides stercoralis* can rarely be found when using the aforementioned microscopy techniques, because its larvae already hatch in the intestine and, hence, the eggs are not passed in the faeces [141]. Despite all these constraints, microscopy is an invaluable tool for diagnostic medical parasitology.

New molecular techniques, especially PCR assays, still need to be validated and further developed in different settings. Disadvantages of current PCR tests are their high costs, risk of contamination, the need for high-technology laboratory equipment and constant electric

Table 5 Diagnostic tests for important viral pathogens that may cause persistent digestive disorders

Infectious pathogen	Diagnostic method				Reference(s)
	Electron microscopy	Cell culture	Immunology	Molecular biology (PCR)	
Viruses					
Adenovirus	Low sensitivity (>10 ⁶ viral particles/ml)	A549-, HEp-2-, HEK-cells	Antigen detection in faecal samples (ELISA, immunochromatography)	PCR	[90]
Astrovirus	Low sensitivity (>10 ⁶ viral particles/ml)	CaCO-2-, LLC-MK2-cells	Antigen detection (ELISA)	RT-PCR	[91]
Bocavirus	-	-	-	PCR	[92]
Coronavirus	-	-	-	RT-PCR	[93]
Cytomegalovirus (CMV)	-	HFF-, MRC-5 cells	• pp65 antigen detection (immunofluorescence)	PCR	[91]
		CMV-immediate early1-pp72-antigen in HFF	• (CMV-specific antibody seroconversion)		
Enterovirus	-	MRC-5-, HEp-2-, Vero-cells	-	RT-PCR	[94]
Human immunodeficiency virus (HIV-1/2)	-	HUT-78-, CEM-MOLT4-cells	• Immunoassay (e.g. 4th generation) • Western Blot	RT-PCR	[95,96]
Norovirus	Sensitivity 10 ⁵ -10 ⁶ viral particles/ml	-	Antigen detection faecal samples (EIA)	RT-PCR	[91]
Parechovirus	-	-	-	RT-PCR	[97]
Rotavirus	Low sensitivity (>10 ⁶ viral particles/ml)	MA104-, CaCO-2-cells	Antigen detection in faecal samples (ELISA), rapid tests (ELISA, immunochromatography)	RT-PCR	[91]
Sapovirus	-	-	-	RT-PCR	[91]

The laboratory techniques are divided into different categories and recommended tests for each pathogen are highlighted. RT-PCR, reverse transcriptase-polymerase chain reaction.




Figure 1 Morphological features of selected intestinal helminth eggs diagnosed using the formalin-ether concentration technique and standard light microscopy: A, *Schistosoma mansoni*; B, *Ascaris lumbricoides*; C, hookworm; D, *Diphylobothrium latum*; E, *Trichuris trichiura*; F, *Capillaria* spp.; G, *Taenia* spp.; H, *Enterobius vermicularis*.

power supply which render their use for routine testing in many developing countries impossible. Indeed, PCR is seldom available in the most affected regions, and its results often do not guide clinicians' decisions, as empiric treatment with albendazole and mebendazole is commonly employed and effective against many helminth species in endemic areas [7]. Due to the variety of intestinal parasites causing digestive disorders, a multiplex real-time PCR targeting a host of various pathogens is much more desirable than individual PCR assays for each parasite, and such multiplex PCRs have been successfully developed and are increasingly used in reference laboratories in industrialised countries [59,83,142]. However, even these multiplex PCRs can only diagnose a defined host of targeted pathogens, while microscopy may sometimes detect unexpected pathogens that would have been missed by other diagnostic methods.

Strongyloides stercoralis

The diagnosis of *S. stercoralis* in human stool samples requires special, often laborious concentration techniques. Most commonly employed are the Baermann funnel and the Koga agar plate [143]. The Baermann method is a concentration technique based on the nematode's hydrophily and thermophily. It provides results within a few hours and is the technique of choice according to the World Gastroenterology Organization [144], but there is some debate whether it is as sensitive as agar plate cultures [86]. Derived from classical charcoal culture assays and its sequel, the so-called Harada-Mori culture, Koga and colleagues developed a special agar plate to detect *S. stercoralis*

and hookworm larvae [145]. The agar plates are stored for 48 hours in a humid chamber and the traces of the helminths can then be seen on the agar and the larvae can easily be collected for microscopic species identification. In contrast to many other helminth infections, where exact species identification often is not necessarily required and clinical symptoms are mild, the recognition of strongyloidiasis and initiation of an effective treatment with ivermectin is essential to prevent potentially life-threatening events due to its ability to cause disseminated hyperinfection in the immunosuppressed population [141,146]. Hence, the aforementioned laborious techniques seem to be justified and a combination of the Baermann funnel and the Koga agar plate method may lead to the most accurate results.

Recently, different PCR assays targeting the helminth's 18S rRNA [87] or 28S rRNA [147] subunit have been developed. First results are promising, but still need further validation in endemic settings.

Schistosoma mansoni*, *S. mekongi*, *S. intercalatum* and *S. japonicum

The microscopic detection of blood fluke eggs in stool specimens still remains the cornerstone of the laboratory diagnosis of intestinal schistosomiasis, as the specificity is high and the costs of equipment are relatively low. However, the sensitivity fluctuates, depending on infection stage and intensity [148]. Hence, concentration methods like an ether-concentration, the Kato-Katz thick smear or the recently developed FLOTAC technique are important tools to increase sensitivity [89].

Moreover, examination of multiple (preferably at least three) stool samples collected on consecutive days is recommended [136,149]. In contrast to other helminth infections, immunological RDTs have been developed for detection of intestinal (*S. mansoni*) and urogenital schistosomiasis (*S. haematobium*). Worm-gut associated glycoproteins, namely circulating cathodic antigen (CCA) and circulating anodic antigen (CAA), can be detected in the serum and the urine of *S. mansoni*-infected individuals using genus-specific monoclonal antibodies [150,151]. Immunochromatographic point-of-care (POC) dipstick or cassette tests for rapid diagnosis of *S. mansoni* via CCA detection in the urine are currently being validated in different epidemiological settings and will potentially become a valuable tool for non-microscopic diagnosis of schistosomiasis in epidemiological studies and clinical practice. Recent studies suggest that the diagnostic accuracy of a single POC-CCA test is considerably more sensitive than a single Kato-Katz thick smear and that a concurrent *S. haematobium* infection does not influence the POC-CCA test results for *S. mansoni* diagnosis, which is an important observation due to the co-endemicity of both blood fluke infections in many tropical areas [152,153]. Hence, antigen RDT assays will likely find their way into clinical practice in the foreseeable future.

PCR assays have been developed and are more sensitive than conventional parasitological and serological methods, but presently, their use is restricted to specialised reference laboratories and research institutions outside endemic areas [154,155].

Viruses

Viral infections commonly cause acute gastroenteritis with the highest burden concentrated in tropical and subtropical regions of the world. Even though these pathogens mainly lead to short-lasting and self-limiting diarrhoeal diseases, they account for considerable morbidity and even mortality, particularly in children [156]. In general, viral infections rarely cause chronic intestinal diseases, but must not be forgotten as potential pathogens that may give rise to persistent diarrhoea and chronic abdominal pain, particularly in HIV-infected individuals or otherwise immunocompromised hosts.

Traditionally, diagnosis of viral gastroenteritis is based on virus isolation by cell culture, electron microscopy and rapid antigen tests (e.g. latex agglutination or EIAs) [157]. Introduction of molecular methods led to an exponential increase in detection rates and the role of difficult-to-culture pathogens became apparent. From a technical point of view, most rapid tests can be done at the bedside, whereas cell culture, electron microscopy and molecular-based methods require laboratories with sophisticated equipment, experienced staff and

appropriate biosafety procedures. This certainly limits the use of the latter methods in resource-constrained settings. Data on sensitivity and specificity of diagnostic tools for virus identification in tropical settings are currently lacking.

Adenovirus

Currently, more than 53 types of adenovirus are recognised which can cause a variety of clinical entities, but gastroenteritis is predominantly caused by types 40 and 41 [158,159]. In infected individuals, viral particles are shed in high concentrations. In general, virus isolation followed by serotyping remains the 'gold' standard for the detection of all serotypes and is possible on different cell lines (Table 5). Importantly, 293-Graham cells should be used for stool samples as adenovirus species F (adenovirus types 40 and 41) can only be cultivated on this cell line. However, virus isolation is rather laborious and time-consuming in the face of urgent requests for diagnosis. Electron microscopy is possible with high specificity, but low sensitivity. As an alternative method that is particularly useful for examination of stool samples, antigen detection assays using EIA or latex agglutination have been developed [160,161]. These assays are rapid, but displayed varying sensitivities and specificities in studies, and hence should be complemented by alternative methods. Molecular methods, in particular real-time PCR, have demonstrated superior performance over conventional methods and are now the cornerstone for diagnosis in most laboratories, but are seldom available in resource-constrained settings.

Astrovirus

Eight serotypes of astrovirus are known. In childhood, astrovirus infection with serotypes 1 and 2 predominate, whereas infection with the other serotypes occurs later in life (>4 years). Prolonged diarrhoea has been associated with astrovirus serotype 3 [162]. In the immunocompetent host, viral shedding occurs for 14–70 hours but may be prolonged in immunosuppressed patients. Virus propagation of astrovirus on CaCO-2 or LLC-MK2 cells remains restricted to expert laboratories and is not recommended for routine diagnostic use. Virus identification by electron microscopy is possible, but appearance of viral particles is not always clear. Recently developed antigen detection kits have proven their suitability and are now widely available for rapid diagnosis. However, sensitivity and specificity of rapid tests in comparison to reverse transcriptase (RT)-PCR have been reported to be comparably low [163]. Real-time RT-PCR is the most sensitive and specific method, but remains restricted to reference laboratories.

Bocavirus

Four different species of human bocavirus (hBoV) have been described thus far [164]. The diagnosis of hBoV infection is almost exclusively based on molecular methods. hBoV has not been isolated by cell culture or in an animal model and rapid antigen tests are currently not available. Serology (e.g. using viral-like particles) has been described and can be used to complement diagnosis [165]. A variety of PCR and real-time PCR assays have been described. However, due to prolonged detection of viral DNA at low copy numbers, qualitative detection of hBoV DNA in gastrointestinal samples is not recommended. There are only few data available for hBoV species 2–4 and the relevance as a true human pathogen is still under debate [164].

Calicivirus

The family *calicivirus* comprises two human-pathogenic genera, the norovirus and sapovirus [166]. For both genera, virus isolation by cell culture is not possible. Electron microscopy is rather insensitive and rarely detects the viruses if there are fewer than 10^6 viral particles/ml of stool suspension.

Norovirus Antigen EIAs have been developed and are commercially available for rapid diagnosis. They proved to be a valuable tool especially in outbreaks, but their sensitivity is limited [167]. A recent study from Brazil reported a sensitivity of 87.9% upon use of a 3rd generation norovirus antigen detection kit [168]. More recently, real-time RT-PCR assays have been described and demonstrated excellent sensitivity and specificity [169]. In-house methods as well as commercial kits are widely available and routinely used.

Sapovirus Specific real-time RT-PCR assays have been developed, but there are no comprehensive data evaluating their diagnostic accuracy. However, there are no diagnostic alternatives because rapid antigen tests are not yet available.

Coronavirus

Five different human pathogenic coronaviruses are known which can cause respiratory and/or to a lesser extent gastrointestinal symptoms in humans. However, the relevance of coronavirus as a true human enteric pathogen is unclear [93,170]. Conventional virus isolation by cell culture can be done on human embryonal tracheal cells. Electron microscopy is possible for stool samples but displays rather low sensitivity. For coronavirus, antigen tests for stool samples are not available. Molecular methods, e.g. real-time RT-PCR assays are the method of choice for a reliable and rapid diagnosis. However,

most in-house methods are restricted to reference laboratories, and hence are not commonly employed around the globe.

Cytomegalovirus

In particular immunosuppressed patients are at risk for cytomegalovirus (CMV) infection, which can affect various organ systems, including the gastrointestinal tract [158,171]. Serology represents the method of choice to differentiate primary from secondary infection. Organ-specific diagnosis (e. g. CMV-associated gastrointestinal disease) requires tissue biopsy samples. In combination with histopathology, isolation of CMV by cell culture is recommended. Detection of CMV-DNA by molecular methods alone is not sufficient.

Enterovirus

Enteroviruses belong to the family *picornaviridae* and comprise enterovirus group A to D [172]. In general, enteroviruses can cause a broad spectrum of different clinical entities. Gastroenteritis caused by coxsackievirus A is mostly seen in children. Virus isolation is possible on a range of different cell lines (Table 5). Virus typing after isolation is traditionally accomplished by virus neutralisation. Of note, enteroviruses may be shed into the stool for prolonged time after clearance of acute infection, thus limiting the significance of such a finding. RT-PCR methods are now widely available for the detection of viral genomes. However, sequence variation among the different enterovirus groups can lower the specificity and PCR-based assays should regularly be updated using latest sequence information. Serological methods for the detection of enterovirus-specific antibodies are exclusively available in reference laboratories and cannot be used for rapid diagnosis.

Parechovirus

Parechoviruses have gained recent interest, but their role in acute gastroenteritis and persistent diarrhoea has yet to be established [173,174]. At the time of writing, 16 parechovirus types have been described. They now represent an own genus within the family *picornaviridae* and real-time RT-PCR is the method of choice for diagnosis in high-income settings [175].

Human rotavirus

Rotavirus infection alone is believed to account for 453,000 deaths annually in children younger than 5 years [5]. In most cases, infection causes acute diarrhoea and vomiting with viral particles being shed in high concentrations. Virus isolation is possible on MA104 or CaCO-2-cells but remains laborious and time-consuming. Antigen detection

by EIA methods is the current standard procedure for the rapid diagnosis of rotavirus infection and widely available for diagnosis as well as surveillance. These assays are able to detect virus particles even if their concentration is below 10^4 particles/ml stool suspension. Molecular methods are also available [176,177].

HIV-associated enteropathy

HIV-associated enteropathy frequently occurs in HIV-infected individuals without access to antiretroviral therapy and is characterised by persistent diarrhoea, weight loss, anorexia, abdominal pain and dysphagia. HIV-associated enteropathy should be diagnosed by obtaining intestinal biopsies via endoscopy with subsequent histological and microbiological examination [178]. Antiretroviral treatment of the HIV infection usually also cures the enteropathy.

Discussion

Persistent digestive disorders are unspecific clinical complaints which are commonly reported by many patients around the world. Gastrointestinal or systemic infections are important causes of such disorders with a broad spectrum of possible pathogens involved, including bacteria, intestinal protozoa, helminths and viruses. Due to the wide range of infectious agents which are often difficult to diagnose, great efforts have to be made to reach satisfactory detection rates and to avoid overlooking of important pathogens. Such a diagnostic work-up should include bacterial stool cultures on different selective media (including MacConkey, sorbitol-MacConkey, Leifson and other agar plate cultures), microscopic examination of unstained (e.g. direct faecal smear, Kato-Katz thick smear and formalin ether-concentration method) and stained microscope slides (acid-fast stains, e.g. Kinyoun technique) for parasite identification, and various pathogen-specific tests such as PCR for viruses and diarrhoeagenic *E. coli* pathotypes, toxin detection kits for *C. difficile* diagnosis, and stool concentration methods for *S. stercoralis* (e.g. Baermann funnel and Koga agar plate). Examination of more than one stool specimen over consecutive days is crucial, because many intestinal pathogens are irregularly shed in the faeces [149]. 'Classical' approaches to persistent diarrhoea often lead to disappointing results with up to 80% of cases in which no causative pathogen can be determined [6].

However, even exhaustive laboratory work-up is prone to a host of limitations and challenges that must be considered and addressed. Firstly, gastrointestinal complaints are often caused by non-infectious causes, and a combination of different clinical signs and symptoms as well as further tests are needed to detect and exclude such non-infectious aetiologies. Secondly, available epidemiological data

regarding the sought infectious pathogens in the tropics are scarce, thus requiring broad diagnostic testing to avoid overlooking of important pathogens. Thirdly, studies should be carried out in different social-ecological settings to assess the influence of cultural, demographic, genetic, geographic, socioeconomic and health system related factors on predominating pathogens. Fourthly, such research must address all pathogen classes and should not be limited to one-dimensional approaches examining either bacteria or parasites only. Fifthly, there are certain issues unique to gastrointestinal diseases which clearly distinguish them from other organ disorders; most importantly, the finding of a given pathogen may not necessarily mean that the patient's complaints are caused by this organism [179]. Bacteria, helminths and intestinal protozoa may often be found as harmless commensals or even beneficial parts of the gastrointestinal flora, and thus such findings may represent coincidence rather than causality [180-182]. This is of particular importance when different potential pathogens are found concurrently in one faecal specimen and the causative one(s) have to be differentiated. Sixthly, even primarily non-intestinal infectious pathogens may cause gastrointestinal symptoms, as has been reported for HIV infection and even malaria in the tropics, where acute or long-lasting diarrhoea may be the only symptom in up to 20% of all observed cases [183,184]. In contrast, patients may as well start to complain about reduced well-being and develop clinical symptoms only some weeks to months after clearance of an intestinal pathogen, as is the case in postinfectious irritable bowel syndrome [185]. Finally, the variety of possible pathogens affecting the gut is so exhaustive that even very sophisticated diagnostic approaches will not be able to detect every pathogen with satisfactory sensitivity and specificity, especially when considering the cost and practical applicability of some specialised techniques that are not currently feasible in most parts of the tropics.

Conclusion

There is a pressing need for research targeting persistent digestive disorders as a coherent clinical problem rather than as a disconnected collection of pathologies. This would allow the elaboration of evidence-based diagnosis-treatment algorithms centred on patients in resource-constrained settings, where data availability is scarce and patient management often driven by experience and local beliefs. This is the overarching goal of the NIDIAG consortium, focusing on digestive disorders as discussed here, as well as on neurological disorders [23] and persistent fever [24]. Additionally, such investigations will optimise the use of existing diagnostic tests and advance the development of new methods, which are ideally able to concurrently detect a broad spectrum of intestinal pathogens with a high sensitivity and specificity, and which are

simple and affordable enough to be performed in low-income countries where prevalences of persistent digestive disorders are generally high. Moreover, the thorough evaluation of reference tests for intestinal pathogens can serve as diagnostic 'gold' standard in the standardisation and validation of easily applicable RDTs, which are highly needed tools in resource-constrained field settings. Finally, such in-depth investigations are not only important for individual patient management, but also for public health policy making (e.g. to assess the efficacy and cost-effectiveness of ongoing preventive chemotherapy control programmes targeting helminthiasis). There is a need for improved diagnostics for persistent digestive disorders in the tropics. It is desirable to conduct a multicentric study to investigate the clinical presentations and respective identified pathogens of large patient cohorts presenting with non-acute gastrointestinal diseases as a first step towards more reliable and evidence-based clinical case management in the tropics.

Competing interest

The authors declare that they have no competing interests.

Authors' contributions

SLB, JV and JU took primary responsibility for the literature search. SLB, JV, SK, MP and JU drafted the manuscript. According to their areas of expertise, the authors critically revised the text chapters (bacteria: SLB, DCW, LvM, CPY and MAM; parasites: SLB, SK, DCW, KP, HM, MS, FM, MAM, LvL, EKN and JU; viruses: MP; clinical aspects: SLB, CPY, JJ, EB and SR). All authors contributed to the manuscript, read and approved the final version.

Financial support

This work is part of the NIDIAG network (Collaborative Project; <http://www.nidiag.org>) supported by the European Commission under the Health Cooperation Work Programme of the 7th Framework Programme (grant agreement no. 260260).

Acknowledgements

We thank Yvette Endriss from the Swiss Tropical and Public Health Institute in Basel, Switzerland, for her help in taking photographs of various helminth eggs for Figure 1.

Author details

¹Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel, Switzerland. ²University of Basel, Basel, Switzerland. ³Institute of Medical Microbiology and Hygiene, University of Saarland Hospital, Homburg/Saar, Germany. ⁴Department of Virology, University of Freiburg, Freiburg, Germany. ⁵Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom. ⁶Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium. ⁷Department of Medical Services and Diagnostic, Swiss Tropical and Public Health Institute, Basel, Switzerland. ⁸J.D. MacLean Centre for Tropical Diseases, and Divisions of Infectious Diseases and Medical Microbiology, McGill University Health Centre, Montreal, Canada. ⁹Department of Clinical Sciences, Institute of Tropical Medicine, Antwerp, Belgium. ¹⁰Institut National de Recherche en Santé Publique, Bamako, Mali. ¹¹Department of Internal Medicine, B P Koirala Institute of Health Sciences, Dharan, Nepal. ¹²Centre for Tropical Medicine, Faculty of Medicine, Gadjah Mada University, Yogyakarta, Indonesia. ¹³Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium. ¹⁴Institut National de Recherche Biomédicale, Kinshasa, Democratic Republic of the Congo. ¹⁵Université de Kinshasa, Kinshasa, Democratic Republic of the Congo. ¹⁶Department of Parasitology, Leiden University Medical Center, Leiden, The Netherlands. ¹⁷Unité de Formation et de Recherche Biosciences, Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire. ¹⁸Département Environnement et Santé, Centre Suisse de

Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire. ¹⁹Division of Humanitarian and International Medicine, Geneva University Hospitals, Geneva, Switzerland.

Received: 17 September 2012 Accepted: 14 January 2013
Published: 24 January 2013

References

1. Bartram J, Cairncross S: **Hygiene, sanitation, and water: forgotten foundations of health.** *PLoS Med* 2010, **7**:e1000367.
2. WHO: *Diarrhoeal disease. Fact sheet no. 330. August 2009.* Geneva: World Health Organization; 2009.
3. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, et al: **Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010.** *Lancet* 2012, **380**:2197–2223.
4. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, Abraham J, Adair T, Aggarwal R, Ahn SY, et al: **Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010.** *Lancet* 2012, **380**:2095–2128.
5. Tate JE, Burton AH, Boschi-Pinto C, Steele AD, Duque J, Parashar UD: **2008 estimate of worldwide rotavirus-associated mortality in children younger than 5 years before the introduction of universal rotavirus vaccination programmes: a systematic review and meta-analysis.** *Lancet Infect Dis* 2012, **12**:136–141.
6. Pawlowski SW, Warren CA, Guerrant R: **Diagnosis and treatment of acute or persistent diarrhea.** *Gastroenterology* 2009, **136**:1874–1886.
7. Utzinger J, Becker SL, Knopp S, Blum J, Neumayr AL, Keiser J, Hatz CF: **Neglected tropical diseases: diagnosis, clinical management, treatment and control.** *Swiss Med Wkly* 2012, **142**:w13727.
8. Pullan RL, Brooker SJ: **The global limits and population at risk of soil-transmitted helminth infections in 2010.** *Parasit Vectors* 2012, **5**:81.
9. Hotez PJ, Molyneux DH, Fenwick A, Ottesen E, Ehrlich Sachs S, Sachs JD: **Incorporating a rapid-impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis, and malaria.** *PLoS Med* 2006, **3**:e102.
10. Utzinger J: **A research and development agenda for the control and elimination of human helminthiasis.** *PLoS Negl Trop Dis* 2012, **6**:e1646.
11. Hotez PJ, Molyneux DH, Fenwick A, Kumaresan J, Ehrlich Sachs S, Sachs JD, Savioli L: **Control of neglected tropical diseases.** *N Engl J Med* 2007, **357**:1018–1027.
12. Hotez P: **Hookworm and poverty.** *Ann N Y Acad Sci* 2008, **1136**:38–44.
13. Hotez PJ, Fenwick A, Savioli L, Molyneux DH: **Rescuing the bottom billion through control of neglected tropical diseases.** *Lancet* 2009, **373**:1570–1575.
14. Conteh L, Engels T, Molyneux DH: **Socioeconomic aspects of neglected tropical diseases.** *Lancet* 2010, **375**:239–247.
15. King CH: **Parasites and poverty: the case of schistosomiasis.** *Acta Trop* 2010, **113**:95–104.
16. Cavalli A, Bamba SI, Traore MN, Boelaert M, Coulibaly Y, Polman K, Pirard M, Van Dormael M: **Interactions between Global Health Initiatives and country health systems: the case of a neglected tropical diseases control program in Mali.** *PLoS Negl Trop Dis* 2010, **4**:e798.
17. Marchal B, Van Dormael M, Pirard M, Cavalli A, Kegels G, Polman K: **Neglected tropical disease (NTD) control in health systems: the interface between programmes and general health services.** *Acta Trop* 2011, **120**(Suppl 1):S177–S185.
18. Amazigo UV, Leak SG, Zoure HG, Njebuome N, Lusamba-Dikassa PS: **Community-driven interventions can revolutionise control of neglected tropical diseases.** *Trends Parasitol* 2012, **28**:231–238.
19. Robays J, Lefevre P, Lutumba P, Lubanza S, Kande Betu Mesu V, Van der Stuyft P, Boelaert M: **Drug toxicity and cost as barriers to community participation in HAT control in the Democratic Republic of Congo.** *Trop Med Int Health* 2007, **12**:290–298.
20. Kibadi K, Boelaert M, Kayinua M, Minuku JB, Muyembe-Tamfum JJ, Portaels F, Lefevre P: **Therapeutic itineraries of patients with ulcerated forms of *Mycobacterium ulcerans* (Buruli ulcer) disease in a rural health zone in the Democratic Republic of Congo.** *Trop Med Int Health* 2009, **14**:1110–1116.

21. Brun R, Blum J, Chappuis F, Burri C: Human African trypanosomiasis. *Lancet* 2010, **375**:148–159.
22. Bix D, de Souza M, Nkengasong JN: Laboratory challenges in the scaling up of HIV, TB, and malaria programs: the interaction of health and laboratory systems, clinical research, and service delivery. *Am J Clin Pathol* 2009, **131**:849–851.
23. Yansouni CP, Bottieau E, Lutumba P, Winkler AS, Lynen L, Büscher P, Jacobs J, Gillet P, Lejon V, Aliro E, et al: Rapid diagnostic tests for neurological infections in Central Africa. *Lancet Infect Dis* 2013, in press.
24. Yansouni CP, Bottieau E, Chappuis F, Phoba MF, Lunguya O, Ifeka BB, Jacobs J: Rapid diagnostic tests for a coordinated approach to fever syndromes in low-resource settings. *Clin Infect Dis* 2012, **55**:610–611.
25. Lacour M, Zunder T, Huber R, Sander A, Daschner F, Frank U: The pathogenetic significance of intestinal *Candida* colonization - a systematic review from an interdisciplinary and environmental medical point of view. *Int J Hyg Environ Health* 2002, **205**:257–268.
26. Abba K, Sinfield R, Hart CA, Garner P: Pathogens associated with persistent diarrhoea in children in low and middle income countries: systematic review. *BMC Infect Dis* 2009, **9**:88.
27. Moore SR: Update on prolonged and persistent diarrhea in children. *Curr Opin Gastroenterol* 2011, **27**:19–23.
28. Pfeiffer ML, Dupont HL, Ochoa TJ: The patient presenting with acute dysentery - a systematic review. *J Infect* 2012, **64**:374–386.
29. Barratt JL, Harkness J, Marriott D, Ellis JT, Stark D: A review of *Dientamoeba fragilis* carriage in humans: several reasons why this organism should be considered in the diagnosis of gastrointestinal illness. *Gut Microbes* 2011, **2**:3–12.
30. Scanlan PD: *Blastocystis*: past pitfalls and future perspectives. *Trends Parasitol* 2012, **28**:327–334.
31. Janda JM, Abbott SL: The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clin Microbiol Rev* 2010, **23**:35–73.
32. Granato PA, Chen L, Holiday I, Rawling RA, Novak-Weekley SM, Quinlan T, Musser KA: Comparison of premier CAMPY enzyme immunoassay (EIA), ProSpecT *Campylobacter* EIA, and ImmunoCard STAT! CAMPY tests with culture for laboratory diagnosis of *Campylobacter* enteric infections. *J Clin Microbiol* 2010, **48**:4022–4027.
33. Bartlett JG, Gerding DN: Clinical recognition and diagnosis of *Clostridium difficile* infection. *Clin Infect Dis* 2008, **46**(Suppl 1):S12–S18.
34. Planche T, Aghaizu A, Holliman R, Riley P, Poloniecki J, Breathnach A, Krishna S: Diagnosis of *Clostridium difficile* infection by toxin detection kits: a systematic review. *Lancet Infect Dis* 2008, **8**:777–784.
35. Lemee L, Dhalluin A, Testelin S, Mattirat MA, Maillard K, Lemeland JF, Pons JL: Multiplex PCR targeting *tpi* (triose phosphate isomerase), *tcdA* (Toxin A), and *tcdB* (Toxin B) genes for toxigenic culture of *Clostridium difficile*. *J Clin Microbiol* 2004, **42**:5710–5714.
36. Kaur P, Chakraborti A, Asea A: Enteroaggregative *Escherichia coli*: an emerging enteric food borne pathogen. *Interdiscip Perspect Infect Dis* 2010, **2010**:254159.
37. Ochoa TJ, Barletta F, Contreras C, Mercado E: New insights into the epidemiology of enteropathogenic *Escherichia coli* infection. *Trans R Soc Trop Med Hyg* 2008, **102**:852–856.
38. Szakal DD, Schneider G, Pal T: A colony blot immune assay to identify enteroinvasive *Escherichia coli* and *Shigella* in stool samples. *Diagn Microbiol Infect Dis* 2003, **45**:165–171.
39. Gould LH, Bopp C, Strockbine N, Atkinson R, Baselski V, Body B, Carey R, Crandall C, Hurd S, Kaplan R, et al: Recommendations for diagnosis of shiga toxin-producing *Escherichia coli* infections by clinical laboratories. *MMWR Recomm Rep* 2009, **58**:1–14.
40. Hermos CR, Janineh M, Han LL, McAdam AJ: Shiga toxin-producing *Escherichia coli* in children: diagnosis and clinical manifestations of O157:H7 and non-O157:H7 infection. *J Clin Microbiol* 2011, **49**:955–959.
41. Nataro JP, Kaper JB: Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev* 1998, **11**:142–201.
42. Guion CE, Ochoa TJ, Walker CM, Barletta F, Cleary TG: Detection of diarrheagenic *Escherichia coli* by use of melting-curve analysis and real-time multiplex PCR. *J Clin Microbiol* 2008, **46**:1752–1757.
43. Wiener RS, Della-Latta P, Schluger NW: Effect of nucleic acid amplification for *Mycobacterium tuberculosis* on clinical decision making in suspected extrapulmonary tuberculosis. *Chest* 2005, **128**:102–107.
44. Giouleme O, Paschos P, Katsaros M, Papalexi F, Karabatsou S, Masmanidou M, Kolioukas D: Intestinal tuberculosis: a diagnostic challenge - case report and review of the literature. *Eur J Gastroenterol Hepatol* 2011, **23**:1074–1077.
45. Bhan MK, Bahl R, Bhatnagar S: Typhoid and paratyphoid fever. *Lancet* 2005, **366**:749–762.
46. Kariuki S: Typhoid fever in sub-Saharan Africa: challenges of diagnosis and management of infections. *J Infect Dev Ctries* 2008, **2**:443–447.
47. Gordon MA: Invasive nontyphoidal *Salmonella* disease: epidemiology, pathogenesis and diagnosis. *Curr Opin Infect Dis* 2011, **24**:484–489.
48. Niyogi SK: Shigellosis. *J Microbiol* 2005, **43**:133–143.
49. Marth T, Schneider T: Whipple disease. *Curr Opin Gastroenterol* 2008, **24**:141–148.
50. Shears P: Recent developments in cholera. *Curr Opin Infect Dis* 2001, **14**:553–558.
51. Neogi SB, Chowdhury N, Asakura M, Hinenoya A, Haldar S, Saidi SM, Kogure K, Lara RJ, Yamasaki S: A highly sensitive and specific multiplex PCR assay for simultaneous detection of *Vibrio cholerae*, *Vibrio parahaemolyticus* and *Vibrio vulnificus*. *Lett Appl Microbiol* 2010, **51**:293–300.
52. Bockemühl J, Roggentin P: Intestinal yersiniosis. Clinical importance, epidemiology, diagnosis, and prevention. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2004, **47**:685–691. (in German).
53. M'ikanatha NM, Dettinger LA, Perry A, Rogers P, Reynolds SM, Nachamkin I: Culturing stool specimens for *Campylobacter* spp., Pennsylvania, USA. *Emerg Infect Dis* 2012, **18**:484–487.
54. Shanholzer CJ, Peterson LR, Olson MN, Gerding DN: Prospective study of gram-stained stool smears in diagnosis of *Clostridium difficile* colitis. *J Clin Microbiol* 1983, **17**:906–908.
55. Schuster FL, Ramirez-Avila L: Current world status of *Balantidium coli*. *Clin Microbiol Rev* 2008, **21**:626–638.
56. Stark D, van Hal S, Marriott D, Ellis J, Harkness J: Irritable bowel syndrome: a review on the role of intestinal protozoa and the importance of their detection and diagnosis. *Int J Parasitol* 2007, **37**:11–20.
57. Stensvold CR, Nielsen HV, Molbak K, Smith HV: Pursuing the clinical significance of *Blastocystis* - diagnostic limitations. *Trends Parasitol* 2009, **25**:23–29.
58. Zhang X, Qiao J, Wu X, Da R, Zhao L, Wei Z: *In vitro* culture of *Blastocystis hominis* in three liquid media and its usefulness in the diagnosis of blastocystosis. *Int J Infect Dis* 2012, **16**:e23–e28.
59. Verweij JJ, Blange RA, Templeton K, Schinkel J, Brien EA, van Rooyen MA, van Lieshout L, Polderman AM: Simultaneous detection of *Entamoeba histolytica*, *Giardia lamblia*, and *Cryptosporidium parvum* in fecal samples by using multiplex real-time PCR. *J Clin Microbiol* 2004, **42**:1220–1223.
60. van Lieshout L, Verweij JJ: Newer diagnostic approaches to intestinal protozoa. *Curr Opin Infect Dis* 2010, **23**:488–493.
61. Caccio SM, Pozio E: Advances in the epidemiology, diagnosis and treatment of cryptosporidiosis. *Expert Rev Anti Infect Ther* 2006, **4**:429–443.
62. Davies AP, Chalmers RM: Cryptosporidiosis. *BMJ* 2009, **339**:b4168.
63. Ortega YR, Sanchez R: Update on *Cyclospora cayentanensis*, a food-borne and waterborne parasite. *Clin Microbiol Rev* 2010, **23**:218–234.
64. Verweij JJ, Mulder B, Poell B, van Middelkoop D, Brien EA, van Lieshout L: Real-time PCR for the detection of *Dientamoeba fragilis* in fecal samples. *Mol Cell Probes* 2007, **21**:400–404.
65. Stark D, Barratt J, Roberts T, Marriott D, Harkness J, Ellis J: Comparison of microscopy, two xenic culture techniques, conventional and real-time PCR for the detection of *Dientamoeba fragilis* in clinical stool samples. *Eur J Clin Microbiol Infect Dis* 2010, **29**:411–416.
66. Qvarnstrom Y, James C, Xayavong M, Holloway BP, Visvesvara GS, Sriram R, da Silva AJ: Comparison of real-time PCR protocols for differential laboratory diagnosis of amoebiasis. *J Clin Microbiol* 2005, **43**:5491–5497.
67. Fotedar R, Stark D, Beebe N, Marriott D, Ellis J, Harkness J: Laboratory diagnostic techniques for *Entamoeba* species. *Clin Microbiol Rev* 2007, **20**:511–532.
68. Clark CG, Diamond LS: Methods for cultivation of luminal parasitic protists of clinical importance. *Clin Microbiol Rev* 2002, **15**:329–341.
69. Kebede A, Verweij JJ, Dorigo-Zetsma W, Sanders E, Messele T, van Lieshout L, Petros B, Polderman T: Overdiagnosis of amoebiasis in the absence of *Entamoeba histolytica* among patients presenting with diarrhoea in Wonji and Akaki, Ethiopia. *Trans R Soc Trop Med Hyg* 2003, **97**:305–307.
70. Kebede A, Verweij JJ, Petros B, Polderman AM: Misleading microscopy in amoebiasis. *Trop Med Int Health* 2004, **9**:651–652.

71. Nagaty IM, Hegazi MM: **Dot-ELISA copro-antigen and direct stool examination in diagnosis of giardiasis patients.** *J Egypt Soc Parasitol* 2007, **37**:641–648.
72. ten Hove RJ, van Lieshout L, Brienen EA, Perez MA, Verweij JJ: **Real-time polymerase chain reaction for detection of *Isospora belli* in stool samples.** *Diagn Microbiol Infect Dis* 2008, **61**:280–283.
73. Murphy SC, Hoogestraat DR, Sengupta DJ, Prentice J, Chakrapani A, Cookson BT: **Molecular diagnosis of cystoisosporiasis using extended-range PCR screening.** *J Mol Diagn* 2011, **13**:359–362.
74. Didier ES, Weiss LM: **Microsporidiosis: not just in AIDS patients.** *Curr Opin Infect Dis* 2011, **24**:490–495.
75. Didier ES, Weiss LM: **Microsporidiosis: current status.** *Curr Opin Infect Dis* 2006, **19**:485–492.
76. Verweij JJ, Ten Hove R, Brienen EA, van Lieshout L: **Multiplex detection of *Enterocytozoon bienersi* and *Encephalitozoon* spp. in fecal samples using real-time PCR.** *Diagn Microbiol Infect Dis* 2007, **57**:163–167.
77. Scholz T, Garcia HH, Kuchta R, Wicht B: **Update on the human broad tapeworm (genus *diphyllobothrium*), including clinical relevance.** *Clin Microbiol Rev* 2009, **22**:146–160.
78. Wicht B, Yanagida T, Scholz T, Ito A, Jimenez JA, Brabec J: **Multiplex PCR for differential identification of broad tapeworms (Cestoda: *Diphyllobothrium*) infecting humans.** *J Clin Microbiol* 2010, **48**:3111–3116.
79. Steinmann P, Cringoli G, Bruschi F, Matthys B, Lohourignon LK, Castagna B, Maurelli MP, Morgogliano ME, Utzinger J, Rinaldi L: **FLOTAC for the diagnosis of *Hymenolepis* spp. infection: proof-of-concept and comparing diagnostic accuracy with other methods.** *Parasitol Res* 2012, **111**:749–754.
80. Brunetti E, White AC Jr: **Cestode infestations: hydatid disease and cysticercosis.** *Infect Dis Clin North Am* 2012, **26**:421–435.
81. Katz N, Chaves A, Pellegrino J: **A simple device for quantitative stool thick-smear technique in schistosomiasis mansoni.** *Rev Inst Med Trop São Paulo* 1972, **14**:397–400.
82. WHO: *Bench aids for the diagnosis of intestinal parasites.* Geneva: World Health Organization; 1994.
83. Basuni M, Muhi J, Othman N, Verweij JJ, Ahmad M, Miswan N, Rahumatullah A, Aziz FA, Zainudin NS, Noordin R: **A pentaplex real-time polymerase chain reaction assay for detection of four species of soil-transmitted helminths.** *AmJ Trop Med Hyg* 2011, **84**:338–343.
84. Soukhathammavong P, Sayasone S, Harimanana AN, Akkhavong A, Thammasack S, Phoumindr N, Choumlivong K, Keoluangkhot V, Phongmany S, Akkhavong K, et al: **Three cases of intestinal capillariasis in Lao People's Democratic Republic.** *AmJ Trop Med Hyg* 2008, **79**:735–738.
85. Attia RA, Tolba ME, Yones DA, Bakir HY, Eldeek HE, Kamel S: ***Capillaria philippinensis* in Upper Egypt: has it become endemic?** *AmJ Trop Med Hyg* 2012, **86**:126–133.
86. Montes M, Sawhney C, Barros N: ***Strongyloides stercoralis*: there but not seen.** *Curr Opin Infect Dis* 2010, **23**:500–504.
87. Verweij JJ, Canales M, Polman K, Ziem J, Brienen EA, Polderman AM, van Lieshout L: **Molecular diagnosis of *Strongyloides stercoralis* in faecal samples using real-time PCR.** *Trans R Soc Trop Med Hyg* 2009, **103**:342–346.
88. Keiser J, Duthaler U, Utzinger J: **Update on the diagnosis and treatment of food-borne trematode infections.** *Curr Opin Infect Dis* 2010, **23**:513–520.
89. Glinz D, Silué KD, Knopp S, Lohourignon LK, Yao KP, Steinmann P, Rinaldi L, Cringoli G, N'Goran EK, Utzinger J: **Comparing diagnostic accuracy of Kato-Katz, Koga agar plate, ether-concentration, and FLOTAC for *Schistosoma mansoni* and soil-transmitted helminths.** *PLoS Negl Trop Dis* 2010, **4**:e754.
90. Henke-Gendo C, Ganzenmueller T, Kluba J, Harste G, Raggub L, Heim A: **Improved quantitative PCR protocols for adenovirus and CMV with an internal inhibition control system and automated nucleic acid isolation.** *J Med Virol* 2012, **84**:890–896.
91. van Maarseveen NM, Wessels E, de Brouwer CS, Vossen AC, Claas EC: **Diagnosis of viral gastroenteritis by simultaneous detection of adenovirus group F, astrovirus, rotavirus group a, norovirus genogroups I and II, and sapovirus in two internally controlled multiplex real-time PCR assays.** *J Clin Virol* 2010, **49**:205–210.
92. Kantola K, Sadeghi M, Antikainen J, Kirveskari J, Delwart E, Hedman K, Soderlund-Venermo M: **Real-time quantitative PCR detection of four human bocaviruses.** *J Clin Microbiol* 2010, **48**:4044–4050.
93. Dare RK, Fry AM, Chittaganpitch M, Sawanpanyalert P, Olsen SJ, Erdman DD: **Human coronavirus infections in rural Thailand: a comprehensive study using real-time reverse-transcription polymerase chain reaction assays.** *J Infect Dis* 2007, **196**:1321–1328.
94. Dierssen U, Rehren F, Henke-Gendo C, Harste G, Heim A: **Rapid routine detection of enterovirus RNA in cerebrospinal fluid by a one-step real-time RT-PCR assay.** *J Clin Virol* 2008, **42**:58–64.
95. Center for Disease Control and Prevention: *Guidelines for appropriate evaluations of HIV testing technologies in Africa.* Available at: http://www.who.int/diagnostics_laboratory/publications/EN_HIVEval_Guide.pdf (accessed online: 26 December 2012).
96. Branson BM: **The future of HIV testing.** *J Acquir Immune Defic Syndr* 2010, **55**(Suppl 2):S102–S105.
97. Baumgarte S, de Souza Luna LK, Grywna K, Panning M, Drexler JF, Karsten C, Huppertz HI, Drosten C: **Prevalence, types, and RNA concentrations of human parechoviruses, including a sixth parechovirus type, in stool samples from patients with acute enteritis.** *J Clin Microbiol* 2008, **46**:242–248.
98. Martiny D, Busson L, Wybo I, El Haj RA, Dediste A, Vandenberg O: **Comparison of the Microflex LT and Vitek MS systems for routine identification of bacteria by matrix-assisted laser desorption/ionization-time of flight mass spectrometry.** *J Clin Microbiol* 2012, **50**:1313–1325.
99. Bessede E, Delcamp A, Sifre E, Buissonniere A, Megraud F: **New methods for detection of campylobacters in stool samples in comparison to culture.** *J Clin Microbiol* 2011, **49**:941–944.
100. Kuhn KG, Falkenhorst G, Ceper T, Dalby T, Ethelberg S, Molbak K, Krogfelt KA: **Detection of antibodies to *Campylobacter* in humans using enzyme-linked immunosorbent assays: a review of the literature.** *Diagn Microbiol Infect Dis* 2012, **74**:113–118.
101. Yaqing H, Wenping Z, Zhiyi Y, Xionghu W, Shouyi Y, Hong Y, Yingchun D, Guifang H: **Detection of human Enterovirus 71 reverse transcription loop-mediated isothermal amplification (RT-LAMP).** *Letts Appl Microbiol* 2012, **54**:233–239.
102. Yamazaki W: **Sensitive and rapid detection of *Campylobacter jejuni* and *Campylobacter coli* using loop-mediated isothermal amplification.** *Methods Mol Biol* 2013, **943**:267–277.
103. Parry CM, Wijedoru L, Arjyal A, Baker S: **The utility of diagnostic tests for enteric fever in endemic locations.** *Expert Rev Anti Infect Ther* 2011, **9**:711–725.
104. Gilman RH, Termini M, Levine MM, Hernandez-Mendoza P, Hornick RB: **Relative efficacy of blood, urine, rectal swab, bone-marrow, and rose-spot cultures for recovery of *Salmonella typhi* in typhoid fever.** *Lancet* 1975, **305**:1211–1213.
105. Baker S, Favorov M, Dougan G: **Searching for the elusive typhoid diagnostic.** *BMC Infect Dis* 2010, **10**:45.
106. Joost I, Speck K, Herrmann M, von Müller L: **Characterisation of *Clostridium difficile* isolates by *slpA* and *tcdC* gene sequencing.** *Int J Antimicrob Agents* 2009, **33**(Suppl 1):S13–S18.
107. Bartlett JG: ***Clostridium difficile*: progress and challenges.** *Ann N Y Acad Sci* 2010, **1213**:62–69.
108. de Jong E, de Jong AS, Bartels CJ, Van Der Rijt-Van Den Biggelaar C, Melchers WJ, Sturm PD: **Clinical and laboratory evaluation of a real-time PCR for *Clostridium difficile* toxin A and B genes.** *Eur J Clin Microbiol Infect Dis* 2012, **31**:2219–2225.
109. Reller ME, Lema CA, Perl TM, Cai M, Ross TL, Speck KA, Carroll KC: **Yield of stool culture with isolate toxin testing versus a two-step algorithm including stool toxin testing for detection of toxigenic *Clostridium difficile*.** *J Clin Microbiol* 2007, **45**:3601–3605.
110. Crobach MJ, Dekkers OM, Wilcox MH, Kuijper EJ: **European Society of Clinical Microbiology and Infectious Diseases (ESCMD): data review and recommendations for diagnosing *Clostridium difficile* infection (CDI).** *Clin Microbiol Infect* 2009, **15**:1053–1066.
111. Robert-Koch-Institut: *Abschließende Darstellung und Bewertung der epidemiologischen Erkenntnisse im EHEC O104:H4 Ausbruch Deutschland 2011.* Berlin: Robert-Koch-Institut, 2011. (in German). Available at: <http://www.rki.de/DE/Content/InfAZ/E/EHEC/EHEC-Abschlussbericht.html> (accessed online: 30 August 2012).
112. Donoghue HD, Holton J: **Intestinal tuberculosis.** *Curr Opin Infect Dis* 2009, **22**:490–496.
113. Jeyanathan M, Boutros-Tadros O, Radhi J, Semret M, Bitton A, Behr MA: **Visualization of *mycobacterium avium* in Crohn's tissue by oil-immersion microscopy.** *Microbes Infect* 2007, **9**:1567–1573.
114. Yu H, Liu Y, Wang Y, Peng L, Li A, Zhang Y: **Clinical, endoscopic and histological differentiations between Crohn's disease and intestinal tuberculosis.** *Digestion* 2012, **85**:202–209.

115. Honscha G, Von Groll A, Valenca M, Ramos DF, Sanchotene K, Scaini CJ, Ribeiro MO, da Silva PE: **The laboratory as a tool to qualify tuberculosis diagnosis.** *Int J Tuberc Lung Dis* 2008, **12**:218–220.
116. Yardley JH, Hendrix TR: **Combined electron and light microscopy in Whipple's disease. Demonstration of "bacillary bodies" in the intestine.** *Bull Johns Hopkins Hosp* 1961, **109**:80–98.
117. Raoult D, Birg ML, La Scola B, Fournier PE, Enea M, Lepidi H, Roux V, Piette JC, Vandenesch F, Vital-Durand D, et al: **Cultivation of the bacillus of Whipple's disease.** *N Engl J Med* 2000, **342**:620–625.
118. Raoult D, Fenollar F, Birg ML: **Culture of *T. whipplei* from the stool of a patient with Whipple's disease.** *N Engl J Med* 2006, **355**:1503–1505.
119. Dobell C: **The discovery of the intestinal protozoa of man.** *Proc R Soc Med* 1920, **13**:1–15.
120. Allen AVH, Ridley DS: **Further observations on the formol-ether concentration technique for faecal parasites.** *J Clin Pathol* 1970, **23**:545–546.
121. Utzinger J, Botero-Kleiven S, Castelli F, Chiodini PL, Edwards H, Kohler N, Gulletta M, Lebbad M, Manser M, Matthys B, et al: **Microscopic diagnosis of sodium acetate-acetic acid-formalin-fixed stool samples for helminths and intestinal protozoa: a comparison among European reference laboratories.** *Clin Microbiol Infect* 2010, **16**:267–273.
122. Becker SL, Lohourignon LK, Speich B, Rinaldi L, Knopp S, N'Goran EK, Cringoli G, Utzinger J: **Comparison of the Flotac-400 dual technique and the formalin-ether concentration technique for diagnosis of human intestinal protozoan infection.** *J Clin Microbiol* 2011, **49**:2183–2190.
123. Sharp SE, Suarez CA, Duran Y, Poppiti RJ: **Evaluation of the Triage Micro Parasite Panel for detection of *Giardia lamblia*, *Entamoeba histolytica*/*Entamoeba dispar*, and *Cryptosporidium parvum* in patient stool specimens.** *J Clin Microbiol* 2001, **39**:332–334.
124. Chalmers RM, Davies AP: **Clinical cryptosporidiosis.** *Exp Parasitol* 2010, **124**:138–146.
125. Diamond LS, Clark CG: **A redescription of *Entamoeba histolytica* Schaudinn, 1903 (Emended Walker, 1911) separating it from *Entamoeba dispar* Brumpt, 1925.** *J Eukaryot Microbiol* 1993, **40**:340–344.
126. Haque R, Neville LM, Hahn P, Petri WA Jr: **Rapid diagnosis of *Entamoeba* infection by using *Entamoeba* and *Entamoeba histolytica* stool antigen detection kits.** *J Clin Microbiol* 1995, **33**:2558–2561.
127. Fotedar R, Stark D, Beebe N, Marriott D, Ellis J, Harkness J: **PCR detection of *Entamoeba histolytica*, *Entamoeba dispar*, and *Entamoeba moshkovskii* in stool samples from Sydney, Australia.** *J Clin Microbiol* 2007, **45**:1035–1037.
128. Ximenez C, Moran P, Rojas L, Valadez A, Gomez A: **Reassessment of the epidemiology of amebiasis: state of the art.** *Infect Genet Evol* 2009, **9**:1023–1032.
129. ten Hove RJ, van Esbroeck M, Vervoort T, van den Ende J, van Lieshout L, Verweij JJ: **Molecular diagnostics of intestinal parasites in returning travellers.** *Eur J Clin Microbiol Infect Dis* 2009, **28**:1045–1053.
130. de Vlas SJ, Gryseels B: **Underestimation of *Schistosoma mansoni* prevalences.** *Parasitol Today* 1992, **8**:274–277.
131. Booth M, Vounatsou P, N'Goran EK, Tanner M, Utzinger J: **The influence of sampling effort and the performance of the Kato-Katz technique in diagnosing *Schistosoma mansoni* and hookworm co-infections in rural Côte d'Ivoire.** *Parasitology* 2003, **127**:525–531.
132. Speich B, Knopp S, Mohammed KA, Khamis IS, Rinaldi L, Cringoli G, Rollinson D, Utzinger J: **Comparative cost assessment of the Kato-Katz and FLOTAC techniques for soil-transmitted helminth diagnosis in epidemiological surveys.** *Parasit Vectors* 2010, **3**:71.
133. García LS, Bruckner DA: *Diagnostic medical parasitology.* Washington D.C: ASM Press; 2001.
134. Tarafder MR, Carabin H, Joseph L, Balolong E Jr, Olveda R, McGarvey ST: **Estimating the sensitivity and specificity of Kato-Katz stool examination technique for detection of hookworms, *Ascaris lumbricoides* and *Trichuris trichiura* infections in humans in the absence of a 'gold standard'.** *Int J Parasitol* 2010, **40**:399–404.
135. Knopp S, Mgeni AF, Khamis IS, Steinmann P, Stothard JR, Rollinson D, Marti H, Utzinger J: **Diagnosis of soil-transmitted helminths in the era of preventive chemotherapy: effect of multiple stool sampling and use of different diagnostic techniques.** *PLoS Negl Trop Dis* 2008, **2**:e331.
136. Berhe N, Medhin G, Erko B, Smith T, Gedamu S, Beredid D, Moore R, Habte E, Redda A, Gebre-Michael T, et al: **Variations in helminth faecal egg counts in Kato-Katz thick smears and their implications in assessing infection status with *Schistosoma mansoni*.** *Acta Trop* 2004, **92**:205–212.
137. Goodman D, Haji HJ, Bickle QD, Stoltzfus RJ, Tielsch JM, Ramsan M, Savioli L, Albonico M: **A comparison of methods for detecting the eggs of *Ascaris*, *Trichuris*, and hookworm in infant stool, and the epidemiology of infection in Zanzibari infants.** *Am J Trop Med Hyg* 2007, **76**:725–731.
138. Steinmann P, Du ZW, Wang LB, Wang XZ, Jiang JY, Li LH, Marti H, Zhou XN, Utzinger J: **Extensive multiparasitism in a village of Yunnan province, People's Republic of China, revealed by a suite of diagnostic methods.** *Am J Trop Med Hyg* 2008, **78**:760–769.
139. Knopp S, Rinaldi L, Khamis IS, Stothard JR, Rollinson D, Maurelli MP, Steinmann P, Marti H, Cringoli G, Utzinger J: **A single FLOTAC is more sensitive than triplicate Kato-Katz for the diagnosis of low-intensity soil-transmitted helminth infections.** *Trans R Soc Trop Med Hyg* 2009, **103**:347–354.
140. Steinmann P, Utzinger J, Du ZW, Zhou XN: **Multiparasitism: a neglected reality on global, regional and local scale.** *Adv Parasitol* 2010, **73**:21–50.
141. Olsen A, van Lieshout L, Marti H, Polderman T, Polman K, Steinmann P, Stothard R, Thybo S, Verweij JJ, Magnussen P: **Strongyloidiasis - the most neglected of the neglected tropical diseases?** *Trans R Soc Trop Med Hyg* 2009, **103**:967–972.
142. Verweij JJ, van Lieshout L: **Intestinal parasitic infections in an industrialized country; a new focus on children with better DNA-based diagnostics.** *Parasitology* 2011, **138**:1492–1498.
143. Becker SL, Sieto B, Silué KD, Adjossan L, Kone S, Hatz C, Kern WV, N'Goran EK, Utzinger J: **Diagnosis, clinical features, and self-reported morbidity of *Strongyloides stercoralis* and hookworm infection in a co-endemic setting.** *PLoS Negl Trop Dis* 2011, **5**:e1292.
144. World Gastroenterology Organization: *WGO practice guideline. Management of strongyloidiasis.* Available at: http://www.worldgastroenterology.org/assets/downloads/en/pdf/guidelines/15_management_strongyloidiasis_en.pdf (accessed online: 1 August 2012).
145. Koga K, Kasuya S, Khamboonruang C, Sukhavat K, Ieda M, Takatsuka N, Kita K, Ohtomo H: **A modified agar plate method for detection of *Strongyloides stercoralis*.** *Am J Trop Med Hyg* 1991, **45**:518–521.
146. Feely NM, Waghorn DJ, Dexter T, Gallen I, Chiodini P: ***Strongyloides stercoralis* hyperinfection: difficulties in diagnosis and treatment.** *Anaesthesia* 2010, **65**:298–301.
147. Kramme S, Nissen N, Soblik H, Erttmann K, Tannich E, Fleischer B, Panning M, Brattig N: **Novel real-time PCR for the universal detection of *Strongyloides* species.** *J Med Microbiol* 2011, **60**:454–458.
148. Engels D, Sinzinkayo E, Gryseels B: **Day-to-day egg count fluctuation in *Schistosoma mansoni* infection and its operational implications.** *Am J Trop Med Hyg* 1996, **54**:319–324.
149. Utzinger J, Booth M, N'Goran EK, Müller I, Tanner M, Lengeler C: **Relative contribution of day-to-day and intra-specimen variation in faecal egg counts of *Schistosoma mansoni* before and after treatment with praziquantel.** *Parasitology* 2001, **122**:537–544.
150. Midzi N, Butterworth AE, Mduluza T, Munyati S, Deelder AM, van Dam GJ: **Use of circulating cathodic antigen strips for the diagnosis of urinary schistosomiasis.** *Trans R Soc Trop Med Hyg* 2009, **103**:45–51.
151. Clerinx J, Van Gompel A: **Schistosomiasis in travellers and migrants.** *Travel Med Infect Dis* 2011, **9**:6–24.
152. Coulibaly JT, Knopp S, N'Guessan NA, Silué KD, Fürst T, Lohourignon LK, Brou JK, N'Gbesso YK, Vounatsou P, N'Goran EK, et al: **Accuracy of urine circulating cathodic antigen (CCA) test for *Schistosoma mansoni* diagnosis in different settings of Côte d'Ivoire.** *PLoS Negl Trop Dis* 2011, **5**:e1384.
153. Shane HL, Verani JR, Abudho B, Montgomery SP, Blackstock AJ, Mwinzi PN, Butler SE, Karanja DM, Secor WE: **Evaluation of urine CCA assays for detection of *Schistosoma mansoni* infection in Western Kenya.** *PLoS Negl Trop Dis* 2011, **5**:e951.
154. ten Hove RJ, Verweij JJ, Vereecken K, Polman K, Dieye L, van Lieshout L: **Multiplex real-time PCR for the detection and quantification of *Schistosoma mansoni* and *S. haematobium* infection in stool samples collected in northern Senegal.** *Trans R Soc Trop Med Hyg* 2008, **102**:179–185.
155. Obeng BB, Aryeetey YA, de Dood CJ, Amoah AS, Larbi IA, Deelder AM, Yazdanbakhsh M, Hartgers FC, Boakye DA, Verweij JJ, et al: **Application of a circulating-cathodic-antigen (CCA) strip test and real-time PCR, in comparison with microscopy, for the detection of *Schistosoma haematobium* in urine samples from Ghana.** *Ann Trop Med Parasitol* 2008, **102**:625–633.
156. Wardlaw T, Salama P, Brocklehurst C, Chopra M, Mason E: **Diarrhoea: why children are still dying and what can be done.** *Lancet* 2010, **375**:870–872.

157. Clark B, McKendrick M: **A review of viral gastroenteritis.** *Curr Opin Infect Dis* 2004, **17**:461–469.
158. Soares CC, Volotao EM, Albuquerque MC, da Silva FM, de Carvalho TR, Nozawa CM, Linhares RE, Santos N: **Prevalence of enteric adenoviruses among children with diarrhea in four Brazilian cities.** *J Clin Virol* 2002, **23**:171–177.
159. Seto D, Chodosh J, Brister JR, Jones MS: **Using the whole-genome sequence to characterize and name human adenoviruses.** *J Virol* 2011, **85**:5701–5702.
160. Grandien M, Pettersson CA, Svensson L, Uhnoo I: **Latex agglutination test for adenovirus diagnosis in diarrheal disease.** *J Med Virol* 1987, **23**:311–316.
161. Gleaves CA, Militoni J, Ashley RL: **An enzyme immunoassay for the direct detection of adenovirus in clinical specimens.** *Diagn Microbiol Infect Dis* 1993, **17**:57–59.
162. Caballero S, Guix S, El-Senousy WM, Calico I, Pinto RM, Bosch A: **Persistent gastroenteritis in children infected with astrovirus: association with serotype-3 strains.** *J Med Virol* 2003, **71**:245–250.
163. Dalton RM, Roman ER, Negredo AA, Wilhelmi ID, Glass RI, Sanchez-Fauquier A: **Astrovirus acute gastroenteritis among children in Madrid, Spain.** *Pediatr Infect Dis J* 2002, **21**:1038–1041.
164. Jartti T, Hedman K, Jartti L, Ruuskanen O, Allander T, Soderlund-Venermo M: **Human bocavirus - the first 5 years.** *Rev Med Virol* 2012, **22**:46–64.
165. Don M, Soderlund-Venermo M, Hedman K, Ruuskanen O, Allander T, Korppi M: **Don't forget serum in the diagnosis of human bocavirus infection.** *J Infect Dis* 2011, **203**:1031–1032. author reply 1032–1033.
166. Green KY: **Caliciviridae: the noroviruses.** In *Fields virology*. 5th edition. Edited by Knipe D, et al. Philadelphia, PA, USA: Lippincott Williams & Wilkins; 2007:949–979.
167. Burton-MacLeod JA, Kane EM, Beard RS, Hadley LA, Glass RI, Ando T: **Evaluation and comparison of two commercial enzyme-linked immunosorbent assay kits for detection of antigenically diverse human noroviruses in stool samples.** *J Clin Microbiol* 2004, **42**:2587–2595.
168. Morillo SG, Luchs A, Cilli A, Ribeiro CD, Calux SJ, Carmona Rde C, Timenetsky Mdo C: **Norovirus 3rd Generation kit: an improvement for rapid diagnosis of sporadic gastroenteritis cases and valuable for outbreak detection.** *J Virol Meth* 2011, **173**:13–16.
169. Gunson RN, Collins TC, Carman WF: **The real-time detection of sapovirus.** *J Clin Virol* 2006, **35**:321–322.
170. Esper F, Ou Z, Huang YT: **Human coronaviruses are uncommon in patients with gastrointestinal illness.** *J Clin Virol* 2010, **48**:131–133.
171. Baroco AL, Oldfield EC: **Gastrointestinal cytomegalovirus disease in the immunocompromised patient.** *Curr Gastroenterol Rep* 2008, **10**:409–416.
172. Nasri D, Bouslama L, Pillet S, Bourlet T, Aouni M, Pozzetto B: **Basic rationale, current methods and future directions for molecular typing of human enterovirus.** *Expert Rev Mol Diagn* 2007, **7**:419–434.
173. Tapia G, Cinek O, Witso E, Kulich M, Rasmussen T, Grinde B, Ronningen KS: **Longitudinal observation of parechovirus in stool samples from Norwegian infants.** *J Med Virol* 2008, **80**:1835–1842.
174. Harvala H, Simmonds P: **Human parechoviruses: biology, epidemiology and clinical significance.** *J Clin Virol* 2009, **45**:1–9.
175. Wolffs PF, Bruggeman CA, van Well GT, van Loo IH: **Replacing traditional diagnostics of fecal viral pathogens by a comprehensive panel of real-time PCRs.** *J Clin Microbiol* 2011, **49**:1926–1931.
176. Pang XL, Lee B, Boroumand N, Leblanc B, Preiksaitis JK, Yu Ip CC: **Increased detection of rotavirus using a real time reverse transcription-polymerase chain reaction (RT-PCR) assay in stool specimens from children with diarrhea.** *J Med Virol* 2004, **72**:496–501.
177. Jothikumar N, Kang G, Hill VR: **Broadly reactive TaqMan assay for real-time RT-PCR detection of rotavirus in clinical and environmental samples.** *J Virol Meth* 2009, **155**:126–131.
178. Bhaijee F, Subramony C, Tang SJ, Pepper DJ: **Human immunodeficiency virus-associated gastrointestinal disease: common endoscopic biopsy diagnoses.** *Patholog Res Int* 2011, **2011**:247923.
179. Lowe AM, Yansouni CP, Behr MA: **Causality and gastrointestinal infections: Koch, Hill, and Crohn's.** *Lancet Infect Dis* 2008, **8**:720–726.
180. Zimmermann MB, Chassard C, Rohner F, N'Goran EK, Nindjin C, Dostal A, Utzinger J, Ghattas H, Lacroix C, Hurrell RF: **The effects of iron fortification on the gut microbiota in African children: a randomized controlled trial in Côte d'Ivoire.** *Am J Clin Nutr* 2010, **92**:1406–1415.
181. Nicholson JK, Holmes E, Kinross J, Burcelin R, Gibson G, Jia W, Pettersson S: **Host-gut microbiota metabolic interactions.** *Science* 2012, **336**:1262–1267.
182. Clemente JC, Ursell LK, Parfrey LW, Knight R: **The impact of the gut microbiota on human health: an integrative view.** *Cell* 2012, **148**:1258–1270.
183. Prasad RN, Virk KJ: **Malaria as a cause of diarrhoea - a review.** *P N G Med J* 1993, **36**:337–341.
184. Reisinger EC, Fritzsche C, Krause R, Krejs GJ: **Diarrhea caused by primarily non-gastrointestinal infections.** *Nat Clin Pract Gastroenterol Hepatol* 2005, **2**:216–222.
185. Spiller R, Garsed K: **Postinfectious irritable bowel syndrome.** *Gastroenterology* 2009, **136**:1979–1988.

doi:10.1186/1471-2334-13-37

Cite this article as: Becker et al.: Persistent digestive disorders in the tropics: causative infectious pathogens and reference diagnostic tests. *BMC Infectious Diseases* 2013 **13**:37.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

